

LETTERS

Discovery of six high-redshift quasars with the Lijiang 2.4 m telescope and the Multiple Mirror Telescope *

Xue-Bing Wu¹, Wen-Wen Zuo¹, Qian Yang¹, Wei-Min Yi², Chen-Wei Yang^{3,4},
Wen-Juan Liu^{3,4}, Peng Jiang^{3,4}, Xin-Wen Shu^{3,4} and Hong-Yan Zhou^{3,4}

¹ Department of Astronomy, Peking University, Beijing 100871, China; wuxb@pku.edu.cn

² National Astronomical Observatories / Yunnan Astronomical Observatory, Chinese Academy of Sciences, Kunming 650011, China

³ Key Laboratory for Research in Galaxies and Cosmology, The University of Science and Technology of China, Chinese Academy of Sciences, Hefei 230026, China

⁴ Polar Research Institute of China, Shanghai 200136, China

Received 2012 June 12; accepted 2012 July 27

Abstract Quasars with redshifts greater than 4 are rare, and can be used to probe the structure and evolution of the early universe. Here we report the discovery of six new quasars with *i*-band magnitudes brighter than 19.5 and redshifts between 2.4 and 4.6 from spectroscopy with the Yunnan Faint Object Spectrograph and Camera (YFOSC) at the Lijiang 2.4 m telescope in February, 2012. These quasars are in the list of $z > 3.6$ quasar candidates selected by using our proposed $J - K/i - Y$ criterion and the photometric redshift estimations from the SDSS optical and UKIDSS near-IR photometric data. Nine candidates were observed by YFOSC, and five among six new quasars were identified as $z > 3.6$ quasars. One of the other three objects was identified as a star and the other two were unidentified due to the lower signal-to-noise ratio of their spectra. This is the first time that $z > 4$ quasars have been discovered using a telescope in China. Thanks to the Chinese Telescope Access Program (TAP), the redshift of 4.6 for one of these quasars was confirmed by the Multiple Mirror Telescope (MMT) Red Channel spectroscopy. The continuum and emission line properties of these six quasars, as well as their central black hole masses and Eddington ratios, were obtained.

Key words: quasars: general — quasars: emission lines — galaxies: active — galaxies: high-redshift

1 INTRODUCTION

The number of known quasars has increased steadily in the past four decades since their discovery in 1963 (Schmidt 1963). In particular, a large number of quasars have been discovered in two large spectroscopic surveys, namely, the Two-degree Field (2dF) survey (Boyle et al. 2000) and the Sloan Digital Sky Survey (SDSS) (York et al. 2000). 2dF mainly selected low redshift ($z < 2.2$) quasar candidates with UV-excess (Smith et al. 2005) and has discovered more than 20 000 quasars (Croom

* Supported by the National Natural Science Foundation of China.

et al. 2004). SDSS adopted a multi-band optical color selection method for quasars mainly by excluding the point sources in the stellar locus of the color-color diagrams (Richards et al. 2002) and has identified more than 120 000 quasars (Schneider et al. 2010). 90% of SDSS quasars have low redshifts ($z < 2.2$), though some dedicated methods were also proposed for finding high redshift quasars ($z > 3.5$) (Fan et al. 2001a,b; Richards et al. 2002).

High-redshift quasars are rare, and those with redshifts greater than 4 represent only 1% of the total quasar population. In the SDSS DR7 quasar catalog (Schneider et al. 2010), only 1248 (392) among 105 783 quasars have redshifts greater than 4 (4.5). Since these $z \sim 4$ quasars existed when the universe was at an age of 1.57 Gyr, they can be used to probe the structure and evolution of the early universe (Smith et al. 1994; Constantin et al. 2002). In particular, the absorption line spectra of these quasars can give valuable information on the nature of the intergalactic medium at high redshifts. However, discovering $z \sim 4$ quasars is a big challenge because they are fainter than the low redshift quasars due to their larger distances. Moreover, the Ly α emission lines for $z \sim 4$ quasars move to the red end of optical spectra, making them hard to distinguish from stars due to similar optical colors. Recently, Wu & Jia (2010) proposed using the $Y - K/g - z$ criterion to select $z < 4$ quasars and using the $J - K/i - Y$ criterion to select $z < 5$ quasars with the SDSS optical and UKIDSS (UKIRT Infrared Deep Sky Survey)¹ near-IR data based on a K -band excess technique (Warren et al. 2000; Hewett et al. 2006; Chiu et al. 2007; Maddox et al. 2008). With these two criteria, we expect to obtain more complete quasar samples than previous ones. Recent optical spectroscopic observations made by the Guo Shou Jing Telescope (LAMOST) and MMT have demonstrated the success of finding the missing quasars with redshifts between 2.2 and 3 using the $Y - K/g - z$ criterion (Wu et al. 2010a,b; Wu et al. 2011). We also hope to discover some $z \sim 4$ quasars with the $J - K/i - Y$ criterion, which is expected to be applicable for selecting the candidates of quasars with redshifts up to 5 (Wu & Jia 2010).

In this letter, we report our discovery of six new high redshift quasars from the spectroscopic observations with the Lijiang 2.4 m telescope and MMT in February, 2012. The successful identifications of these high redshift quasars further demonstrate the effectiveness of using our newly proposed criteria for discovering the missing quasars including high-redshift ones.

2 TARGET SELECTION AND SPECTROSCOPIC OBSERVATIONS

Richards et al. (2009) presented a catalog of about one million quasar candidates selected from the SDSS DR6 photometric data using Bayesian methods. Photometric redshifts for these candidates were also provided based on the SDSS *ugriz* magnitudes. From this catalog we selected all unidentified candidates with photometric redshift greater than 3.6, photometric redshift probability larger than 0.6 and *i*-band magnitude brighter than 19.5. Then we cross-matched them with the UKIDSS Large Area Survey (LAS) DR7 catalog using a positional offset of 3 arcsec to find the closest counterparts. From this sample with both SDSS *ugriz* data and UKIDSS *YJHK* data, we adopted our $J - K/i - Y$ criterion (Wu & Jia 2010), namely, $J - K > 0.45(i - Y) + 0.48$ (where *YJK* are the Vega magnitudes and *i* is the AB magnitude), to make further selection of $z \sim 4$ quasar candidates. We also used our own program to estimate the photometric redshifts of these candidates with SDSS and UKIDSS 9-band photometric data (Wu & Jia 2010; Wu et al. 2004), and excluded the sources whose photometric redshifts, estimated from the 5-band SDSS photometric data in Richards et al. (2009), are inconsistent with ours. After performing these procedures, we obtained a final list of about 20 high-redshift ($z > 3.6$) quasar candidates.

The spectroscopic observations were carried out on February 26–28, 2012, with the Yunnan Faint Object Spectrograph and Camera (YFOSC) instrument of the Lijiang 2.4 m telescope in

¹ The UKIDSS project is defined in Lawrence et al. (2007). UKIDSS uses the UKIRT Wide Field Camera (WFCAM; Casali et al. 2007) and a photometric system described in Hewett et al. (2006). The pipeline processing and science archive are described in Hambly et al. (2008).

Table 1 Parameters of Nine Objects Observed by YFOSC

Name (SDSS J)	Date	Exposure (s)	<i>u</i>	<i>g</i>	<i>r</i>	<i>i</i>	<i>z</i>	<i>Y</i>	<i>J</i>	<i>H</i>	<i>K</i>	Result
075733.86+190403.1	2012-02-26	2700	21.37	20.40	19.45	19.00	18.53	17.88	17.02	16.63	15.68	low S/N
085203.84+020437.7	2012-02-27	6000	21.67	20.99	19.69	19.09	18.66	17.82	17.47	16.66	15.72	low S/N
092740.04-023347.5	2012-02-28	4200	21.01	20.50	19.55	19.08	18.86	18.42	18.01	17.11	16.42	G star
093345.70-020439.5	2012-02-27	3600	23.77	20.72	19.55	19.47	19.41	19.21	18.66	18.46	17.88	quasar
095023.74+004419.7	2012-02-27	6600	23.97	20.97	19.75	19.48	19.36	18.83	18.40	17.76	17.25	quasar
113816.85+045023.6	2012-02-27	6000	21.26	20.70	19.67	19.27	19.06	18.30	17.96	16.90	16.46	quasar
120312.63-001118.8	2012-02-28	5400	25.38	22.34	20.29	19.14	18.95	18.32	18.00	17.19	16.64	quasar
125052.11+074919.6	2012-02-26	5400	23.56	20.08	18.75	18.63	18.43	18.14	17.35	16.81	16.15	quasar
145115.89+015843.3	2012-02-27	4800	23.72	20.42	19.30	19.23	19.09	18.61	18.02	17.56	16.92	quasar

The *ugriz* magnitudes are given in the AB system and the *YJHK* magnitudes are given in the Vega system.




Fig. 1 The YFOSC spectra of six new quasars. From the left to right, the red dashed lines mark the wavelengths of Ly α , SiIV and CIV emission lines at the estimated redshift for five $z > 3.6$ quasars, while for SDSS J113816.85+045023.6 they mark the wavelengths of CIV and CIII] (*color online*).

Yunnan Astronomical Observatory. Due to the cloudy weather, nine candidates were observed with YFOSC using a low resolution grism with central wavelength around 6500 Å, spectral resolving power of 870, and a long slit with a width of 2.5". The typical seeing was around 2". In Table 1 we summarize the details of the observations for these nine candidates. Six of them were identified as quasars, one as a G-type star and two as unidentified due to the low signal-to-noise ratios of their spectra.

The spectra of six new quasars, after the flat-field correction and both wavelength and flux calibrations, are shown in Figure 1. The strongest emission lines for five $z > 3.6$ quasars are Ly α lines, while for SDSS J113816.85+045023.6 the strongest line around 6400 Å is CIII]. The redshifts for these quasars are the average values given mostly by the Ly α and CIV lines for the five $z > 3.6$ quasars and by the CIII] and CIV lines for SDSS J113816.85+045023.6.




Fig. 2 The MMT Red Channel spectrum of SDSS J120312.63–001118.8. The average redshift estimated from Ly α and SiIV (1400 Å) emission lines is 4.601. The red line refers to the smoothed spectrum with a bin size of 8 Å (*color online*).

Thanks to the Chinese Telescope Access Program², SDSS J120312.63–001118.8 was also observed with the Red Channel Spectrograph on the MMT 6.5m telescope³ at Mt. Hopkins, Arizona, USA on 2012 Feb. 29, with a wavelength coverage of 5100–8600 Å and a spectral resolution of 1.6 Å. It was observed twice, with exposures of 10 min and 15 min respectively. The spectrum was processed using the IRAF Echellette package and is shown in Figure 2. The average redshift estimated from Ly α and SiIV (1400 Å) emission lines is 4.601 ± 0.008 , consistent with the result ($z = 4.592 \pm 0.048$) obtained from the YFOSC observation.

3 SPECTRAL ANALYSES AND PROPERTIES OF SIX HIGH-REDSHIFT QUASARS

The redshift corrected rest-frame YFOSC spectra of six quasars are first corrected for the Galactic extinction using the extinction map of Schlegel et al. (1998). They are then fitted with an IDL code MPFIT (Markwardt 2009). We fit the spectra with the pseudo-continuum model consisting of the featureless nonstellar continuum and FeII emissions. The continuum is assumed to be a power-law, so two free parameters (the amplitude and the slope) are required. The UV FeII template (Vestergaard & Wilkes 2001; Tsuzuki et al. 2006) is convolved with a velocity dispersion and shifted with a velocity, assuming that the line widths of FeII lines are the same as those of emission lines in the corresponding wavelength range. During the fitting, the amplitude and slope of the power-law continuum and the amplitude, velocity shift and broadening width of the FeII emission are set to be free parameters. The initial value of the power-law continuum is obtained by fitting a simple power law to the data in the chosen windows, which are free from emission-line contamination. The initial value of broadening width of the FeII emissions is set to be the line width of the strong emission line CIV. Then we use the pseudo-continuum model to fit a set of continuum windows with strong FeII emissions but no other emission lines, as mentioned in Hu et al. (2008), slightly adjusted interactively for each individual spectrum in order to avoid broad absorption features or extended wings of emission lines.

After constructing the pseudo-continuum, the CIV line should be fitted with two Gaussians, one for the narrow component and the other for the broad component. However, except SDSS J095023.74+004419.7, the spectra of the other five quasars have low signal-to-noise ratio, so we used only one Gaussian to fit the CIV emission line. Absorption features are evident in the spec-

² <http://info.bao.ac.cn/tap/>

³ Observation reported here was obtained at the MMT Observatory, a joint facility of the University of Arizona and the Smithsonian Institution.

Table 2 Spectral Parameters and Black Hole Masses of Six New Quasars

Name (SDSS J)	Redshift	FWHM (CIV) (km s ⁻¹)	log (L_{1350}) (erg s ⁻¹)	log (M_{BH}) (M_{\odot})	log (L_{bol}) (erg s ⁻¹)	log R_{EDD}
093345.70-020439.5	3.701±0.011	6749	46.457	9.621	47.122	-0.588
095023.74+004419.7	3.967±0.035	4500	46.300	9.185	46.964	-0.310
113816.85+045023.6	2.368±0.011	5144	46.000	9.118	46.664	-0.543
120312.63-001118.8	4.592±0.048	4865	46.431	9.323	47.096	-0.316
125052.11+074919.6	3.748±0.030	5424	46.805	9.615	47.469	-0.235
145115.89+015843.3	3.940±0.006	4507	45.682	8.859	46.346	-0.602

tra of four quasars, and one more negative Gaussian was added in the fitting. We measure the Full Width at Half Maximum of the CIV line (FWHM_{CIV}), with luminosity at 1350 Å (L_{1350}) from the spectra (except for SDSS J113816.65+004419.7, where 1350 Å is not within the wavelength coverage, and we estimate the luminosity at 1500 Å instead). The black hole mass is estimated based on FWHM_{CIV} and L_{1350} with equation (7) in Vestergaard & Peterson (2006) (see also Kong et al. 2006). Using a scaling between L_{1350} and bolometric luminosity L_{bol} , $L_{\text{bol}} = 4.62L_{1350}$, we estimated the bolometric luminosity for the six quasars. Based on the estimated black hole mass and bolometric luminosity, we also estimated their Eddington ratios (R_{EDD}). The results are summarized in Table 2. Although the uncertainties of these values are probably quite large due to the low spectral quality and the unusual properties of CIV, we noticed that the overall properties of these six quasars are consistent with those of typical SDSS quasars at high redshifts (Shen et al. 2011).

4 DISCUSSION

A complete quasar sample is crucial for studying the large scale structure of the universe. The current available quasar samples are mostly biased towards low redshifts ($z < 2.2$) and more efforts are needed to find quasars at high redshifts. Wu & Jia (2010) proposed to obtain a larger, more complete quasar sample with redshifts up to five by combining the $J - K/i - Y$ criterion with the $Y - K/g - z$ criterion to select quasar candidates. Some recent optical spectroscopic observations have demonstrated the success of finding the missing quasars with redshifts between 2.2 and 3 using the $Y - K/g - z$ criterion (Wu et al. 2010a,b; Wu et al. 2011). Our discovery of six high redshift quasars (five with $z > 3.6$) from the spectroscopic observations with the Lijiang 2.4 m telescope and MMT further demonstrates the effectiveness of using the $J - K/i - Y$ criterion for discovering quasars with redshifts up to five. Moreover, the identification of five quasars with $z > 3.6$ from nine candidates with photometric redshifts larger than 3.6 also confirms the robustness of the photometric redshifts estimated by the SDSS and UKIDSS photometric data. We noticed that two among our five $z > 3.6$ new quasars do not meet the SDSS *gri* or *riz* selection criterion for $z > 3.6$ quasars (Fan et al. 2001a,b; Richards et al. 2002), which suggests that about 40% of such quasars may be missed in the SDSS spectroscopic survey. This obviously needs to be confirmed by future observations of a large sample of $z > 3.6$ quasars.

Our identification of a $z = 4.6$ quasar demonstrates that $z > 4$ quasars can be identified with the 2-meter size telescopes in China. We hope more high redshift quasars will be discovered by the future LAMOST quasar survey (Wu & LAMOST Extragalactic Survey Team 2011), which is aiming at discovering 0.3–0.4 million quasars from 1 million quasar candidates with $i < 20.5$, by taking advantage of 4000 fibers and the 5 degree field of view of LAMOST (Su et al. 1998; Zhao et al. 2012). The new quasar selection criteria, such as those based on SDSS, UKIDSS and the Wide-field Infrared Survey Explorer (WISE; Wright et al. 2010) data (Wu et al. 2012), will be applied for selecting quasar candidates in the LAMOST quasar survey. This will hopefully provide the largest quasar sample in the next five years for further studies of AGN physics, large scale structure and cosmology.

Acknowledgements We thank the referee for a constructive report and Jianguo Wang, Cheng Hu and Zhaoyu Chen for great helps on the spectral analysis. This work was supported by the National Natural Science Foundation of China (Grant No. 11033001). We acknowledge the use of the Lijiang 2.4 m telescope and the MMT 6.5 m telescope, as well as the archive data from SDSS and UKIDSS. This research uses data obtained through the Telescope Access Program (TAP), which is funded by the National Astronomical Observatories, Chinese Academy of Sciences, and the Special Fund for Astronomy from the Ministry of Finance. Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is <http://www.sdss.org/>.

References

- Boyle, B. J., Shanks, T., Croom, S. M., et al. 2000, *MNRAS*, 317, 1014
 Casali, M., Adamson, A., Alves de Oliveira, C., et al. 2007, *A&A*, 467, 777
 Chiu, K., Richards, G. T., Hewett, P. C., & Maddox, N. 2007, *MNRAS*, 375, 1180
 Constantin, A., Shields, J. C., Hamann, F., et al. 2002, *ApJ*, 565, 50
 Croom, S. M., Smith, R. J., Boyle, B. J., et al. 2004, *MNRAS*, 349, 1397
 Fan, X., Strauss, M. A., Schneider, D. P., et al. 2001a, *AJ*, 121, 54
 Fan, X., Narayanan, V. K., Lupton, R. H., et al. 2001b, *AJ*, 122, 2833
 Hambly, N. C., Collins, R. S., Cross, N. J. G., et al. 2008, *MNRAS*, 384, 637
 Hewett, P. C., Warren, S. J., Leggett, S. K., & Hodgkin, S. T. 2006, *MNRAS*, 367, 454
 Hu, C., Wang, J.-M., Ho, L. C., et al. 2008, *ApJ*, 687, 78
 Kong, M.-Z., Wu, X.-B., Wang, R., & Han, J.-L. 2006, *ChJAA (Chin. J. Astron. Astrophys.)*, 6, 396
 Lawrence, A., Warren, S. J., Almaini, O., et al. 2007, *MNRAS*, 379, 1599
 Maddox, N., Hewett, P. C., Warren, S. J., & Croom, S. M. 2008, *MNRAS*, 386, 1605
 Markwardt, C. B. 2009, in *ASP Conference Series*, 411, *Astronomical Data Analysis Software and Systems XVIII*, eds. D. A. Bohlender, D. Durand, & P. Dowler (San Francisco: ASP), 251
 Richards, G. T., Fan, X., Newberg, H. J., et al. 2002, *AJ*, 123, 2945
 Richards, G. T., Myers, A. D., Gray, A. G., et al. 2009, *ApJS*, 180, 67
 Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
 Schmidt, M. 1963, *Nature*, 197, 1040
 Schneider, D. P., Richards, G. T., Hall, P. B., et al. 2010, *AJ*, 139, 2360
 Shen, Y., Richards, G. T., Strauss, M. A., et al. 2011, *ApJS*, 194, 45
 Smith, J. D., Djorgovski, S., Thompson, D., et al. 1994, *AJ*, 108, 1147
 Smith, R. J., Croom, S. M., Boyle, B. J., et al. 2005, *MNRAS*, 359, 57
 Su, D. Q., Cui, X., Wang, Y., & Yao, Z. 1998, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 3352, ed. L. M. Stepp, 76
 Tsuzuki, Y., Kawara, K., Yoshii, Y., et al. 2006, *ApJ*, 650, 57
 Vestergaard, M., & Peterson, B. M. 2006, *ApJ*, 641, 689
 Vestergaard, M., & Wilkes, B. J. 2001, *ApJS*, 134, 1
 Warren, S. J., Hewett, P. C., & Foltz, C. B. 2000, *MNRAS*, 312, 827
 Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868
 Wu, X.-B., Zhang, W., & Zhou, X. 2004, *ChJAA (Chin. J. Astron. Astrophys.)*, 4, 17
 Wu, X.-B., Chen, Z.-Y., Jia, Z.-D., et al. 2010a, *RAA (Research in Astronomy and Astrophysics)*, 10, 737
 Wu, X.-B., Jia, Z.-D., Chen, Z.-Y., et al. 2010b, *RAA (Research in Astronomy and Astrophysics)*, 10, 745
 Wu, X.-B., & LAMOST Extragalactic Survey Team 2011, *arXiv:1111.0738*
 Wu, X.-B., Wang, R., Schmidt, K. B., et al. 2011, *AJ*, 142, 78
 Wu, X.-B., Hao, G., Jia, Z., et al. 2012, *AJ*, 144, 49
 Wu, X.-B., & Jia, Z. 2010, *MNRAS*, 406, 1583
 York, D. G., Adelman, J., Anderson, J. E., Jr., et al. 2000, *AJ*, 120, 1579
 Zhao, G., Zhao, Y.-H., Chu, Y.-Q., et al. 2012, *RAA (Research in Astronomy and Astrophysics)*, 12, 723