Z Draconis with two companions in a 2:1 mean-motion resonance

Jin-Zhao Yuan¹, Hakan Volkan Şenavcı², and Sheng-Bang Qian³

¹ Department of Physics, Shanxi Normal University, Linfen 041004, China; yuanjz@sxnu.edu.cn

² Department of Astronomy and Space Sciences, Ankara University, Faculty of Science, TR-06100 Tandoğan-Ankara, Turkey

³ Yunnan Observatories, Chinese Academy of Sciences, 650011 Kunming, China

Received 2015 October 3; accepted 2015 November 28

Abstract All available mid-eclipse times of the eclipsing binary Z Draconis are analyzed, and three sets of cyclic variations with periods of 20.1, 29.96 and 59.88 yr are found. The low-amplitude variations with a period of 20.1 yr may be attributed to the unavoidable slight imperfection in the double-Keplerian model, which gives periods of 29.96 and 59.88 yr. Interestingly, the Z Draconis system is close to a 2:1 mean-motion resonance, or a 6:3:2 mean-motion resonance if the 20.1 yr period really exists. We also find that the best solutions tend to give the minimum eccentricities. Based on Kepler's third law, the outermost companion has a minimum mass of $\sim 0.77~M_{\odot}$, whereas the middle companion is an M dwarf star with a mass of $\sim 0.40~M_{\odot}$, suggesting that Z Draconis is a general N-body system.

Key words: binaries: close - stars: individual (Z Draconis) - methods: numerical

1 INTRODUCTION

The oscillations in the mid-eclipse times of eclipsing binaries are usually explained as light-travel time (LTT) effect and magnetic activity cycles (Applegate 1992; Yuan & Qian 2007). In the LTT model, a companion revolves around the eclipsing pair. The line-of-sight distance between the eclipsing pair and the barycenter of the whole system, d, varies with a strict period equal to the orbital period of the companion. After dividing by the speed of light, c, we obtain the O - C value, d/c. Obviously, the multiperiodic variations in the eclipse times of an eclipsing binary provide us important constraints on the orbital characteristics of this multi-companion system, which is usually comprised of an eclipsing binary and multiple sub-stellar objects or planets. In the magnetic activity mechanism, the gravitational or magnetic forces change as the active component goes through a magnetic activity cycle, producing quasi-periodic variations in the eclipse times (Beuermann et al. 2012).

Z Draconis (BD+73°533 = HIP 57348, $V_{\text{max}} = 10.67$ mag) was first found to be an Algol-type binary (hereafter Z Dra AB) by Ceraski (1903). Due to its high declination and brightness, a large number of photometric data were obtained by small telescopes. The first radial velocity curve for the primary component was obtained by Struve (1947). Based on the radial velocity curve and the *BVRI* light curves obtained with a 0.25 Schmidt-Cassegrain telescope, Terrell (2006) carried out a photometric-spectroscopic analysis. The solutions indicated that Z Dra is a semi-detached binary with masses of $1.47 M_{\odot}$ for the primary component and $0.43 M_{\odot}$ for the secondary component. Terrell (2006) also pointed out that the mass of the primary is significantly lower than expected for an A5V star, but consistent with the B-V color of 0.45 mag. Dugan (1915) conducted a detailed period study of the system and found that the mid-eclipse times show two sinusoidal variations with periods of 10.7 and 26.8 yr, while Rafert (1982) found only one cyclic period of 20.3 yr. However, many mid-eclipse times have been obtained in the past 32 yr. Therefore, it is necessary to reanalyze the behavior of the change in the observed period.

In this paper, the O-C data are derived from all available mid-eclipse times in Section 2, where we also present several new data. In Section 3, we apply the fitting procedures described in Yuan & Şenavci (2014, hereafter Paper I). In Section 4, we test the Keplerian model, and find that most of the best-fit elements are valid. Finally, we summarize our results and give our conclusions in Section 5.

2 ECLIPSE-TIMING VARIATIONS

We carried out CCD observations of Z Dra in 2014 April and 2015 February using the 40-cm Schmidt-Cassegrain telescope at the Ankara University Kreiken Observatory in Turkey (AUKR-T40), and the 60-cm Cassegrain telescope at Yunnan Observatories (YNAO-60) in China. The exposure times we adopted in 2014 April are 60 s, 30 s, 20 s, and 15 s in B, V, R and I bands, respectively. The exposure times in 2015 February were 80 s in the V band and 50 s in the R band. The comparison and check stars were BD+72°545 ($\alpha_{J2000.0} = 11^{h}45^{m}56.1^{s}$, $\delta_{J2000.0} = 72°05'44.5''$) and GSC 4395–201 ($\alpha_{J2000.0} = 11^{h}43^{m}21.5^{s}$, $\delta_{J2000.0} = 72°06'34.2''$), respectively. The data reduction was performed by using the aperture photometry package IRAF¹ (bias subtraction, flat-field division). Extinction corrections were ignored as the comparison star is very close to the variable. We fit the transit center of the eclipse by using the technique of Kwee & van Woerden (1956). In total, three new mid-eclipse times are obtained and are listed in Table 1.

The Lichtenknecker Database of the BAV² and the O-C Gateway Database³ list all of the available mid-eclipse times of Z Dra in the literature. In addition, 15 mid-eclipse times between 1928 and 1949 were obtained by Kreiner et al. (2001) and kindly sent to us (via private communication). Three visual and photographic times (HJD 2415787.7856, 2451728.4900 and 2453209.4470) are discarded due to their large deviation from the O - C curve. In total, we have collected 820 mid-eclipse times, which have a time span of 125 yr. All of the data are plotted in Figure 1.

Most mid-eclipse times were published without uncertainties. Therefore, a probable uncertainty of $\sigma = \pm 0.0003$ d is assumed for the photoelectric and CCD data, and ± 0.005 d for the photographic, plate and visual data. Considering that the CCD times obtained simultaneously in different filters may differ from each other by as much as ± 0.0003 d, the uncertainty of ± 0.0003 d is adopted if the mid-eclipse time was obtained in a single filter with uncertainty less than ± 0.0003 d. Eventually, all high-precision (i.e., $\sigma < 0.001$ d) data are spread over the last 16 yr, and most low-precision (i.e., $\sigma > 0.001$ d) data over the remaining time.

Since the Heliocentric Julian Dates (HJDs) in the Coordinated Universal Time (UTC) system are not uniform, all eclipse times after 1950 have been converted to Barycentric Julian Dates (BJDs) in the Barycentric Dynamical Time (TDB) system using the UTC2BJD³ procedure provided by Eastman et al. (2010). For the eclipse times before 1950, the relation between Universal Time (UT) and Terrestrial Time (TT) given by Duffett-Smith & Zwart (2011) is adopted for for a visual conversion, producing additional uncertainties of a few seconds, which are much smaller than their assumed uncertainty of 0.005 d (i.e., 432 s).

Based on the eclipse times between 2011 and 2014, a new linear ephemeris

$$\operatorname{Min} I = \operatorname{HJD2456775.4604} + 1^{d}.35745406 \times E \quad (1)$$

is obtained for future observations. In this paper, the eclipse-timing residuals, O-C, are computed with respect



Fig. 1 The O - C diagram of Z Dra based on the 820 original data. Note that the error bars are smaller than the squares for the high-precision data in the last 4000 cycles.

to the linear ephemeris given by Kreiner et al. (2001),

$$Min I = BJD2443499.7305 + 1^{d}.35743190 \times E, \quad (2)$$

where E denotes the cycle number. The O - C data are displayed in Figure 1.

3 DATA ANALYSIS AND LTT MODELS

Usually, there is mass transfer between two components in an Algol-type binary, and the observed period should increase or decrease, suggesting that the O - C curve has a parabolic trend. It is obvious that a single parabola cannot describe the O - C curve very well, implying that an additional periodic model may be required. Since the data are sampled unevenly with different uncertainties, it is inappropriate to use the parabolic model and the periodic model in turn. If the residuals of a best fit are used for another fit, then one would obtain a best-fit solution different from that given by a combination of both models. Therefore, a quadratic plus sinusoidal model

$$O - C = T_O(E) - T_C(E) = C_0 + C_1 \times E + C_2 \times E^2 + A \sin(2\pi t/P_3) + B \cos(2\pi t/P_3)$$
(3)

is used to calculate the generalized Lomb-Scargle (GLS) periodogram, which is plotted in Figure 2(a). In Equation (3), A, B and $C_{0,1,2}$ are free coefficients. In the GLS periodogram, the power peaks at 18.8, 20.5 and 56.0 yr. As pointed out by Zechmeister & Kürster (2009), the GLS periodogram can give a good initial guess for the best Keplerian period with only a slight frequency shift.

Then, we simultaneously use a second-order polynomial and one LTT term to fit the O - C values

$$O - C = T_O(E) - T_C(E) = C_0 + C_1 \times E + C_2 \times E^2 + \tau_3,$$
(4)

¹ IRAF is developed by the National Optical Astronomy Observatory, which are operated by the Association of Universities for Research in Astronomy, Inc., under contract with the National Science Foundation.

² http://www.bav-astro.de/index.php?sprache=en

³ http://var.astro.cz/ocgate/

³ http://astroutils.astronomy.ohio-state.edu/time/

Table 1 Several New Mid-eclipse Times of Z Dra

HJD (UTC)	BJD (TE	DB)	Error	Mir	n. Filte	r Origin
24000	+00	240000	0+	(d)			
56775	.4605	56775.46	127	± 0.0002	Ι	В	AUKR-T40
56775	.4605	56775.46	127	± 0.0002	Ι	V	AUKR-T40
56775	.4602	56775.46	097	± 0.0002	Ι	R	AUKR-T40
56775	.4603	56775.46	107	± 0.0002	Ι	Ι	AUKR-T40
57063	.2409	57063.24	167	± 0.0002	Ι	V	YNAO-60
57063	.2410	57063.24	177	± 0.0002	Ι	R	YNAO-60
57071	.3858	57071.38	657	± 0.0002	Ι	V	YNAO-60
57071	.3860	57071.38	677	± 0.0002	Ι	R	YNAO-60

where the LTT term τ_3 is derived from Keplerian orbits (Irwin 1952), and can be expressed as

$$\tau_3 = \frac{a_3 \sin i_3}{c} \Big[\frac{1 - e_3^2}{1 + e_3 \cos \nu_3} \sin(\nu_3 + \omega_3) + e_3 \sin \omega_3 \Big].$$
(5)

In Equation (5), $a_3 \sin i_3$ is the projected semimajor axis of the eclipsing pair around the barycenter of the triple system (i_3 is the orbital inclination of the companion with respect to the tangent plane of the sky). e_3 is the eccentricity and ω_3 is the argument of the periastron measured from the ascending node in the tangent plane of the sky. ν_3 is the true anomaly, which is related to the mean anomaly $M = 2\pi(t - T_3)/P_3$, where T_3 and P_3 are the time of the periastron passage and orbital period, respectively.

For fixed e_3 , T_3 and P_3 , ν_3 can be computed for all mid-eclipse times. Then, we fit the O - C data with Equation (4), and get the goodness-of-fit statistic, χ^2 , which is the weighted sum of the squared difference between the O - C values y_i and the model values $y(t_i)$ at eclipse times t_i

$$\chi^{2} = \sum_{i=1}^{N} \left[\frac{y_{i} - y(t_{i})}{\sigma_{i}} \right]^{2} = W \sum_{i=1}^{N} w_{i} \left[y_{i} - y(t_{i}) \right]^{2}, \quad (6)$$

where

$$w_i = \frac{1}{W} \frac{1}{\sigma_i^2},\tag{7}$$

and

$$W = \sum_{j=1}^{N} \frac{1}{\sigma_j^2}.$$
 (8)

In Equation (6), σ_i is the uncertainty associated with O-C of the data point y_i , and N is the number of data points.

Stepping through e_3 and T_3 , we obtain the local χ^2 minimum for a fixed P_3 , i.e., $\chi^2(P_3)$. Since $\sum_{i=1}^N w_i = 1$, $\sqrt{\chi^2(P_3)/W}$ can be regarded as the weighted root mean square (rms) scatter around the best fit for a fixed P_3 (Marsh et al. 2014). After searching P_3 , the global chi-square minimum, χ^2_{global} , can be found. Some local χ^2 minima at $P_3 > 100$ yr give the companion with a mass more than $200 M_{\odot}$, so these are ruled out by us. Normalized by χ^2_{global} , we obtain a power spectrum (Zechmeister & Kürster 2009; Cumming et al. 1999; Cumming 2004),

$$p(P_3) \equiv \frac{\chi_0^2 - \chi^2(P_3)}{\chi_{\text{global}}^2},$$
 (9)

where the constant χ_0^2 is the best-fit value of the χ^2 statistic for a fit of a parabola to the data. Figure 2(b) shows the one-dimensional Keplerian periodogram as well as the best-fit eccentricity $e(P_3)$. If the minimum rms scatter, $\sqrt{\chi_{\text{global}}^2/W}$, is taken as noise in the power spectrum and $\sqrt{(\chi_0^2 - \chi^2)/W}$ as a signal, the $\sqrt{p(P_3)}$ would be the signal-to-noise ratio.

Both one-dimensional periodograms show an extremely significant periodicity at ~60 yr, suggesting a companion with a period of ~60 yr (hereafter, referred to as Z Dra (AB)C). The power also peaks at P = ~30 yr, suggesting another companion with an orbital period of ~30 yr (hereafter, referred to as Z Dra (AB)D). The companion is in/around a 2:1 mean-motion resonance (MMR) with Z Dra (AB)C. The ~20 yr periodicity reported by Rafert (1982) is obvious in Figure 2(a) and 2(b). If the ~20 yr signal exists, the eclipsing binary has a third companion (hereafter, Z Dra (AB)E). It is interesting that Z Dra (AB)C, D and E are in 6:3:2 MMRs. Furthermore, the best-fit eccentricity, $e(P_3)$, reaches local minimum values near $P_3 = ~60$, ~30 and ~20 yr.

We also note that, due to the short time coverage, the power increases continuously from ~ 80 yr, but always remains below the ~ 60 yr peak. Although the power at long periods $(P_3 > 80 \text{ yr})$ is still large, the best-fit solutions at long periods give an eccentricity larger than 0.70. Such a large eccentricity is physically unlikely. A large eccentricity often implies a large gravitational perturbation from other companions. The statistic $p(P_3)(N-8)/4$ follows Fisher's F distribution with 4 and N - 8 degrees of freedom (Bevington & Robinson 1992). Integrating the distribution function and multiplying it by the number of independent frequencies gives a false alarm probability (FAP) less than 10^{-30} for the three peaks (Cumming et al. 1999; Cumming 2004). In fact, the FAP values should be derived from a suitable model, but the one-companion model is not suitable for the Z Dra system (see below).

The best fits corresponding to the 60 yr periodicity are plotted in Figure 3(a), and listed in the second column



Fig. 2 The GLS periodogram (a) and Keplerian periodogram (b) of Z Dra. The dashed vertical lines mark three peaks in the Keplerian periodogram. The red line represents the best-fit eccentricity corresponding to $\chi^2(P_3)$.

(Solution 1) of Table 2. For safety, we also use a thirdorder polynomial instead of the second-order polynomial in Equation (4), and obtain Solution 2, which is shown in Figure 3(b). As shown in the bottom panels of Figure 3(a) and 3(b), the residuals at \sim BJD2446000 reach as large as 0.02 d, which are much larger than their uncertainties. It seems that the residuals show cyclic variation with a period of \sim 30 yr.

To determine further whether there are two periodicities in the O - C data, we also use a second-order polynomial plus two-LTT ephemeris to fit the O - C data. We search for the best period in 40–90 yr with one LTT term, and the other LTT term in 10–40 yr. The linearized Keplerian fitting method (Beuermann et al. 2012; Paper I), which is very similar to the method of the one-dimensional periodogram above, is used to calculate a two-dimensional periodogram. The least-squares fit to the 820 data involves thirteen free parameters, three for the second-order polynomial in the ephemeris, and five orbital elements (P_k, e_k , ω_k, T_k and $a_k \sin i_k/c$) for each companion. If all the parameters are free, then the number of degrees of freedom (DOF) is therefore 807. The constraints on the two orbital periods are shown in Figure 4(a). The χ^2 contour levels of 1.05, 1.2, 1.5, 2.0, 3.0, 4.0 and 5.0 have been normalized by division of the global chi-square minimum, χ^2_{global} . In addition, the best-fit eccentricities of Z Dra (AB)C and D are plotted in Figure 4(b) and Figure 4(c), respectively. In the two-dimensional periodogram, the global χ^2 minimum at ($P_4 \simeq 60$ yr, $P_3 \simeq 30$ yr) confirms Z Dra (AB)C and D, and the local χ^2 minimum at ($P_4 \simeq 60$ yr, $P_3 \simeq 20$ yr) reveals Z Dra (AB)C and E. Both χ^2 minima lie close to the points of the e_3 minima and also the e_4 minima.

Based on the best solution in the two-dimensional periodogram, the Levenberg-Marquardt fitting algorithm (Markwardt 2009) is adopted to search for improved solutions. The improved fits are plotted in Figure 5(a) and 5(b). The corresponding parameters and χ^2 are listed in the fourth and fifth columns (i.e., Solutions 3 and 4) of Table 2. After the parabolic trend is removed, the residuals are displayed in Figure 6, where two sets of periodic variations can be seen more clearly. Compared to Solution 1, the reduced $\chi^2_{\nu} = 2.6$ ($\chi^2 = 2118.3$ for 807 DOF) in Solution 3 is greatly improved, but is still unacceptable (Bradt 2004). The large χ^2_{ν} is due to the large uncertainties in the old O - C data before BJD2415700 (i.e., E < -20500), and perhaps a third set of cyclic variations in the residuals.



Fig.3 The one-companion fit to the eclipse-timing variations of Z Dra. (a) The overplotted solid line denotes the best fit with Equation (3), and the dashed line only represents the second-order polynomial in the ephemeris. The residuals of the best fit are displayed in the lower panel. (b) The same as figure (a) but a third-order polynomial is adopted.

Parameter	Solution 1	Solution 2	Solution 3	Solution 4
C_0 (d)	-0.0197 ± 0.0001	-0.0211 ± 0.0010	-0.0025 ± 0.0007	-0.0146 ± 0.0009
$C_1 \; (\times 10^{-6} \; \mathrm{d})$	$4.80 {\pm} 0.01$	$4.44 {\pm} 0.03$	$4.49 {\pm} 0.02$	$4.88 {\pm} 0.07$
$C_2 \; (\times 10^{-10} \; \mathrm{d})$	$4.35 {\pm} 0.01$	$4.80 {\pm} 0.20$	$3.94{\pm}0.01$	3.16 ± 0.03
$C_3 \; (\times 10^{-15} \; \mathrm{d})$		$-3.85 {\pm} 0.25$		-5.06 ± 0.30
P_4 (yr)			$29.81 {\pm} 0.08$	$29.05 {\pm} 0.08$
T_4 (BJD)			2400464 ± 164	$2403688 {\pm} 513$
e_4			$0.43 {\pm} 0.01$	$0.11 {\pm} 0.03$
$a_4 \sin i_4$ (au)			$2.14 {\pm} 0.03$	$1.92 {\pm} 0.09$
ω_4 (deg)			285.6 ± 3.9	$83.1 {\pm} 26.0$
$m_4 \ (M_{\odot}, i_4 = 90^{\circ})$			$0.39 {\pm} 0.03$	$0.33 {\pm} 0.04$
$A_4 \; (au, i_4 = 90^\circ)$			12.74 ± 0.3	12.3 ± 0.2
P_3 (yr)	$57.49 {\pm} 0.27$	59.11 ± 0.15	59.41±0.12	58.07 ± 0.12
T_3 (BJD)	2411634 ± 77	2410320 ± 127	2401400 ± 109	$2412114{\pm}101$
e_3	$0.42 {\pm} 0.01$	$0.41 {\pm} 0.01$	$0.62 {\pm} 0.02$	$0.56 {\pm} 0.01$
$a_3 \sin i_3$ (au)	$5.52 {\pm} 0.06$	$5.79 {\pm} 0.07$	$6.05 {\pm} 0.07$	$5.61 {\pm} 0.06$
ω_3 (deg)	232.9 ± 1.2	226.4 ± 1.1	$76.8 {\pm} 4.0$	240.5 ± 1.6
$m_3 (M_{\odot}, i_3 = 90^{\circ})$	$0.70 {\pm} 0.01$	$0.73 {\pm} 0.01$	$0.77 {\pm} 0.02$	$0.77 {\pm} 0.03$
$A_3 (au, i_3 = 90^\circ)$	20.5 ± 0.3	$20.9 {\pm} 0.4$	22.3 ± 0.3	$21.9 {\pm} 0.2$
χ^2	6221.1	6110.6	2118.3	2005.8



Fig.4 (a) Two-dimensional Keplerian periodogram derived from a second-order polynomial plus two-LTT model. The χ^2 contours of 1.05, 1.2, 1.5, 2.0, 3.0, 4.0 and 5.0 have been normalized by division of the global χ^2 minimum. (b) The best-fit eccentricity (e_3) of Z Dra (AB)C as a function of (P_3 , P_4). The darker the color is, the smaller the eccentricity is. (c) The same as figure (b) but for e_4 .

As shown in Figure 5(a) and 5(b), the LTT signal of Z Dra (AB)E can be seen in the residuals of the twocompanion fit. A further fit reveals that Z Dra (AB)E has an orbital period of $P_5 = \sim 20.1$ yr and a mass of $\sim 0.2 M_{\odot}$. Z Dra (AB)E produces a cyclic O - C variation with a semi-amplitude of $a_5 \sin i_5 = \sim 0.8$ au, which is much smaller than $a_3 \sin i_3$. In such a case, it is also possible that such a small signal arises from an unavoidable slight imperfection in the double-Keplerian model (see below).

4 TESTS OF THE SO-CALLED KEPLERIAN MODEL

Based on an assumed inclination for one companion, its mass (m_k) can be estimated from the following mass functions

$$\frac{(m_4 \sin i_4)^3}{(m_b + m_4)^2} = \frac{4\pi^2}{GP_4{}^2} \times (a_4 \sin i_4)^3, \tag{10}$$

$$\frac{(m_3 \sin i_3)^3}{(m_b + m_4 + m_3)^2} = \frac{4\pi^2}{GP_3^2} \times (a_3 \sin i_3)^3, \quad (11)$$

where G is the Newtonian gravitational constant. For simplicity, the central eclipsing binary is treated as a single

object $(m_{\rm b})$ with a mass equal to the sum of the masses of both components. In the case of Z Dra, $m_{\rm b} = 1.90 M_{\odot}$ (Terrell 2006). It is important to keep in mind that the m_4 and m_3 derived in this way are just approximate masses since the mass functions are derived from Kepler's third law. If the orbital inclinations of both companions are 90.0°, then the outer companion Z Dra (AB)C has a minimum mass of $\sim 0.8~M_{\odot}$, whereas the inner companion Z Dra (AB)D is an M dwarf with a mass of $\sim 0.4 M_{\odot}$. It is obvious that Z Dra is a general N-body system. Given m_4 and m_3 , we can calculate the semimajor axes of the two companions by the equations, $A_4 = a_4 \cdot (m_b + m_4)/m_4$ and $A_3 = a_3 \cdot (m_b + m_3 + m_4)/m_3$. The minimum A_4 is about 420 times larger than the separation between Z Dra A and B ($6.38R_{\odot} = 0.030$ au), suggesting that the central eclipsing pair can be treated as a single object.

Assuming that Z Dra (AB)C and D revolve around Z Dra AB in coplanar Keplerian orbits with $i_3 = i_4 = 90.0^\circ$, the centripetal force (F_c) of Z Dra (AB)D from the eclipsing pair is comparable with the gravitational perturbation (F_p) from the outer companion, Z Dra (AB)C. The "relative perturbation," F_p/F_c , is calculated on a 130 yr timescale (BJD 2410000 – 2457482). In the process of



Fig.5 The two-companion fit to the eclipse-timing variations of Z Dra when a second-order polynomial trend (a) or a third-order polynomial trend (b) is considered. The residuals of the best fit are displayed in the lower panel of each figure. The overplotted solid line denotes the best fit with a polynomial plus two-LTT ephemeris and the dashed line only represents the polynomial in the ephemeris.



Fig. 6 The same as Fig. 1 but subtracted by the parabolic trend given by Solution 3.

calculation, we track the coordinates of the three objects. Then, the forces of gravity are derived from their separations and masses.

For Solution 3 or 4, the result reveals that F_p/F_c peaks at ~ 0.25 with a mean value of ~ 0.09 (see Fig. 7). The gravitational perturbation can decrease if the errors of the orbital parameters, especially $\omega_{3,4}$ and $T_{3,4}$, are considered. Although mutually tilted orbits can also reduce the gravitational perturbation, the mutual perturbation between the two companions cannot be neglected. The Keplerian formula serves only as a convenient mathematical description of the O - C data.



Fig.7 The ratio of the gravitational perturbations from the outer companion (Z Dra (AB)C) to centripetal forces from the central eclipsing pair (Z Dra A and B), both of which act on Z Dra (AB)D in opposition to each other. The solid line is derived from Solution 3, and the dashed line is from Solution 4.

Generally, the Newtonian LTT signals derived from N-body simulations differ more or less from those given by the double-Keplerian model (Marsh et al. 2014; Goździewski et al. 2012, 2015). If we are only interested in the orbital periods of two companions, the LTT value caused by the outer companion, $(O-C)_3$, can still be fitted by the LTT model given by Equation (4). In this case, the best-fit parameters have no physical meaning except for the orbital period and the projected semimajor axis. (Strictly, $a_3 \sin i_3$ is half of the width of the orbit in the line-of-sight direction.) Comparing with a sinusoidal model with three free parameters (i.e., $A\sin(Bt + C)$), the LTT model has five free parameters, and can be used to generate a larger variety of more complex O - C curves (see Fig. 8). There must be a Keplerian O - C curve whose shape is the most similar to the true $(O - C)_3$. The best-fit Keplerian curve may differ slightly from the true $(O - C)_3$. Note that the true O - C value is equal to $(O - C)_3$ plus $(O - C)_4$ if the parabolic trend is neglected. Such slight deviations would have some influence on a second fit to $(O - C)_4$, which is caused by the inner companion. In Solutions 3 and 4, $a_4 \sin i_4$ is about one third of $a_3 \sin i_3$, suggesting that the influence on the second fit is also at a low level. Therefore, the result of the 2:1 MMR is valid, and the masses of Z Dra (AB)C and D are approximate. As for the low-amplitude $(a_5 \sin i_5 = 0.8 \text{ au})$ variation, it may arise from a slight imperfection in the double-Keplerian model.

Finally, we would like to remind the reader that the best-fit eccentricities are not exactly equal to the orbital eccentricities. Actually, the observed eccentricity results from the true orbital eccentricity and the deviation of the angular velocity from that predicted by Keplerian motion. The deviation of the angular velocity arises from the gravitational perturbation from other companions, and therefore should be small since the gravitational perturbation should be small in a stable system. On the other hand, the true orbital eccentricity should also be small since a companion with small orbital eccentricity often experiences a weak gravitational perturbation from other companions. These may be the reason why the χ^2 minima lie close to points of $e_{3,4}$ minima.

5 DISCUSSION AND CONCLUSIONS

Detailed O - C analyses of Z Dra are performed by using all of the available mid-eclipse times in the literature as well as three new mid-eclipse times obtained in this paper. The O - C diagram shows a quadratic or cubic trend. A companion with orbital period more than twice as long as the time window of observation can produce a quadratic/cubic O - C curve, which is actually a section of a cyclic O-C curve. However, the quadratic/cubic trend is often explained by mass transfer between two components. The quadratic trend in Solution 3 represents an observed period increase with a rate of $dP/dt = 2.1 \times 10^{-7} d \text{ yr}^{-1}$, which is a typical value for many contact binary stars (see e.g., Qian 2001, 2003; Qian et al. 2008). The cubic trend in Solution 4 suggests that the observed period increases at a decreasing rate. The mass transfer from the secondary component to the primary will cease in 197 years. Then, the mass will be transferred from the primary component to the secondary one. Compared with binary evolutionary timescales, a timescale of a few hundred years is negligibly short. The mass transfer rate in the eclipsing pair should change little or remain constant over a few hundred years, suggesting a quadratic trend rather than a cubic trend. Furthermore, the cubic model does not have a significant advantage over the quadratic model. The best-fit cubic trend in Figure 5(b) is close to the quadratic trend in Figure 5(a). Similar periodicities and χ^2 are obtained in Solutions 3 and 4.

We have searched the O - C data for periodicities. The O - C data show two or more sets of cyclic variations with periods of 59.4 and 29.8, and possible ~ 20.1 and > 80 yr, suggesting there are two or more companions around the eclipsing binary. Although we cannot ascertain the exact number of companions, there must be more than one companion. If only one companion revolves around



Fig.8 All kinds of O - C curves derived from by the Keplerian model given by Equation (5). Different colors refer to different eccentricities, and different line styles and thicknesses denote the O - C curves with different ω values. The semi-amplitudes of all O - C curves are normalized to unity. The orbital phase is proportional to time, and has been shifted so that phase zero corresponds to the *BJD* time of the O - C maximum.



Fig.9 Schematic positions of the eclipsing binary (m_b) and its two companions $(m_{3,4})$ with respect to the tangent plane of the sky (Paper I). $i_{3,4}$ denote the orbital inclinations of $m_{3,4}$, respectively. $r_{3,4}$ refer to Jacobian coordinates of $m_{3,4}$. Note that the subscripts '3' and '4' are assigned to the outer and inner companions, respectively.

the eclipsing binary, and therefore moves in a Keplerian orbit, then the single-Keplerian model would fit the O - Cdata very well. However, Figure 3(a) and (b) shows that the single-Keplerian model fails. The two-dimensional periodogram reveals that the companions Z Dra (AB)C and D with periods of 59.4 and 29.8 yr are the most likely combination. As for the long period (> 80 yr), Figure 4(b) and (c) shows any long-period companion has large e_3 and e_4 , which lie far from the points representing the e_3 and e_4 minima, respectively. Such large eccentricities are physically unlikely.

Although magnetic activity can explain the cyclic variations in the O - C diagram (Applegate 1992; Yuan & Qian 2007), they are unlikely to produce two/three sets of variations with commensurate periods. The more plausible reason for such variations is the reflex motion of the eclipsing pair induced by two/three companions in a 2:1 or 6:3:2 MMR. In Paper I, two companions were found to be in near 3:1 MMR orbits around the eclipsing binary SW Lac with periods of 27.0 and 82.6 yr. Both Z Dra and SW Lac have the most numerous mid-eclipsing times, which show complex variations. Perhaps, MMRs are common in such N-body systems.

More than 160 planetary systems have been confirmed so far. About 30% of them are close to MMRs, particularly near the first order MMRs of 2:1 and 3:2 (Zhang et al. 2014). Furthermore, Beuermann et al. (2013) found that two planetary companions are in near 2:1 MMR orbits around the eclipsing binary NN Ser. However, in contrast to these planetary systems, Z Dra and SW Lac are general three-body systems if the central eclipsing binary is treated as a single object. Our discoveries will help us understand the orbital properties of such three-body systems.

In this paper, we have checked the mutual perturbations between Z Dra (AB)C and D, but are unable to test the dynamical stability for three reasons: (1) The so-called double-Keplerian model only gives a convenient approximation to the O - C curve, but it cannot provide exact orbits and correct initial conditions (such as coordinates, velocities and masses) for the N-body system. (2) The errors of the orbital parameters should be considered in our dynamic simulations. (3) The inclinations $(i_3 \text{ and } i_4)$ and the angle between their ascending nodes in the sky plane (θ) are unknown. The orbital angular configuration of the outer companion relative to the inner one is determined by i_3 , i_4 and θ (see Fig. 9). It seems extremely difficult to test the dynamical stability since stable configurations are likely to be confined to tiny regions of parameter space for the general three-body system and all initial conditions must be very accurate in the dynamical analyses.

We note that the N-body model was used to fit the LTT data of HU Aqr (Goździewski et al. 2012, 2015) and NN Ser (Marsh et al. 2014), both of which host two circumbinary planets. In this model, synthetic LTT signals at all epoches are determined through numerical N-body integration, and then compared to the true LTT signals. Based on the reduced χ^2_{ν} and dynamical stability, one can find the best solution. In the case of Z Dra, the masses of Z Dra (AB)C and D are relatively large. We have to fit the LTT data with the masses, velocities and coordinates as free parameters, by integrating the equations of motion (K. Goádziewski, private communication). For a general three-body system, seven free parameters are needed for each companion, giving 17 free parameters in the model. Besides the computational challenges, a possible drawback of this model is that an undiscovered companion can make the N-body integration meaningless. If the orbital period of the companion is more than twice as long as the time window of observation, then the LTT signals caused by the companion would show a parabolic/cubic trend in the time window rather than periodic variation. In this case, one would miss the third companion, and therefore its dynamical perturbation. The long-period companion, however, has little influence on any analytic model including a second-order or third-order polynomial, such as the thirdorder polynomial plus double-Keplerian ephemeris.

As shown in Figures 2 and 4, none of the periodograms can provide tight constraints on the periodicity of > 70 yr. This is mainly attributed to the low precision of the old data and the short time coverage of the O - C data. Therefore, we encourage follow-up observations of this system to obtain more mid-eclipse times covering as long a baseline as possible.

Acknowledgements We would like to thank J. M. Kreiner for his data and the anonymous referee for some constructive suggestions. This research has also made use of the Lichtenknecker-Database of the BAV, operated by the Bundesdeutsche Arbeitsgemeinschaft für Veränderliche Sterne e.V. (BAV). The computations were carried out at National Supercomputer Center in Tianjin, and the calculations were performed on TianHe-1(A). This work is supported by the National Natural Science Foundation of China (NSFC, No. U1231121) and the research fund of Ankara University (BAP) (No. 13B4240006).

References

- Applegate, J. H. 1992, ApJ, 385, 621
- Beuermann, K., Dreizler, S., & Hessman, F. V. 2013, A&A, 555, A133
- Beuermann, K., Dreizler, S., Hessman, F. V., & Deller, J. 2012, A&A, 543, A138
- Bevington, P. R., & Robinson, D. K. 1992, Data Reduction and Error Analysis for the Physical Sciences (2nd ed.; New York: McGraw-Hill)
- Bradt, H. 2004, Astronomy Methods: A Physical Approach to Astronomical Observations (Cambridge: Cambridge Univ. Press)
- Ceraski, W. 1903, Astronomische Nachrichten, 161, 159
- Cumming, A. 2004, MNRAS, 354, 1165
- Cumming, A., Marcy, G. W., & Butler, R. P. 1999, ApJ, 526, 890
- Duffett-Smith, P., & Zwart, J. 2011, Practical Astronomy with your Calculator or Spreadsheet (Cambridge: Cambridge Univ. Press)
- Dugan, R. S. 1915, MNRAS, 75, 702
- Eastman, J., Siverd, R., & Gaudi, B. S. 2010, PASP, 122, 935
- Goździewski, K., Nasiroglu, I., Słowikowska, A., et al. 2012, MNRAS, 425, 930
- Goździewski, K., Słowikowska, A., Dimitrov, D., et al. 2015, MNRAS, 448, 1118
- Irwin, J. B. 1952, ApJ, 116, 211
- Kreiner, J. M., Kim, C.-H., & Nha, I.-S. 2001, An Atlas of O-C Diagrams of Eclipsing Binary Stars (Cracow, Poland: Wydawnictwo Naukowe Akademii Pedagogicznej)
- Kwee, K. K., & van Woerden, H. 1956, Bull. Astron. Inst. Netherlands, 12, 327
- Markwardt, C. B. 2009, in Astronomical Society of the Pacific Conference Series, 411, Astronomical Data Analysis Software and Systems XVIII, eds. D. A. Bohlender, D. Durand, & P. Dowler, 251
- Marsh, T. R., Parsons, S. G., Bours, M. C. P., et al. 2014, MNRAS, 437, 475
- Qian, S. 2001, MNRAS, 328, 914
- Qian, S. 2003, MNRAS, 342, 1260
- Qian, S.-B., He, J.-J., Liu, L., Zhu, L.-Y., & Liao, W. P. 2008, AJ, 136, 2493
- Rafert, J. B. 1982, PASP, 94, 485
- Struve, O. 1947, ApJ, 106, 92
- Terrell, D. 2006, Information Bulletin on Variable Stars, 5742, 1
- Yuan, J., & Şenavci, H. V. 2014, MNRAS, 439, 878
- Yuan, J., & Qian, S. 2007, ApJ, 669, L93
- Zechmeister, M., & Kürster, M. 2009, A&A, 496, 577
- Zhang, X., Li, H., Li, S., & Lin, D. N. C. 2014, ApJ, 789, L23