

Global photometric analysis of Galactic HII regions

Anastasiia Topchieva¹, Dmitri Wiebe¹ and Maria S. Kirsanova^{1,2}

¹ Institute of Astronomy, Russian Academy of Sciences, Moscow 119017, Russia; stasyat@inasan.ru

² Ural Federal University, 19 Mira Str., Ekaterinburg, Russia

Received 2017 November 29; accepted 2018 January 12

Abstract Total infrared fluxes are estimated for 99 HII regions around massive stars. The following wavebands have been used for the analysis: 8 and 24 μm , based on data from the *Spitzer Space Telescope* (IRAC and MIPS, respectively); 70, 160, 250, 350 and 500 μm , based on data from the *Herschel Space Observatory* (PACS and SPIRE). The estimated fluxes are used to evaluate the polycyclic aromatic hydrocarbon (PAH) mass fraction (q_{PAH}) and the intensity of ultraviolet emission in the studied objects. It is shown that the PAH mass fraction, q_{PAH} , is much lower in these objects than the average Galactic value, implying effective destruction of aromatic particles in HII regions. Estimated radiation field intensities (U) are close to those derived for extragalactic HII complexes. Color indices $[F_{24}/F_8]$, $[F_{70}/F_{24}]$, $[F_{160}/F_{24}]$ and $[F_{160}/F_{70}]$ are compared to criteria proposed to distinguish between regions of ionized hydrogen and planetary nebulae. Also, we relate our results to analogous color indices for extragalactic complexes of ionized hydrogen.

Key words: ISM: bubbles — (ISM:) HII regions — ISM: lines and bands

1 INTRODUCTION

The amount of new data on the infrared (IR) radiation in our Galaxy grows steadily. Thanks to results, which have been obtained with the *Spitzer Space Telescope*, we now have the opportunity to study objects which had been known previously as ring nebulae and are now widely referred to as IR bubbles (Churchwell et al. 2006, 2007). Their formation is presumably related to the action of massive hot stars on interstellar material (van Buren & McCray 1988). Specifically, it is believed that a bubble appears around an O-B type star, which ionizes surrounding gas and forms an expanding shell due to hot gas pressure and/or powerful stellar wind.

Observations tend to support this picture. Deharveng et al. (2010) classified 86% of objects from the Churchwell et al. (2006) catalog as HII regions. Anderson et al. (2014) created a catalog which includes more than 8000 Galactic HII regions and HII region candidates, using a specific morphology in the mid-IR band as a selection criterion. We should also mention results from the Milky Way project (Simpson et al. 2012) and a catalog by Makai et al. (2017), based on observations from the *Spitzer* and *WISE* space telescopes.

The number of detected objects increases each year, and finally we can lay out a solid basis for statistical and theoretical studies. A statistical analysis of HII regions (see e.g. Anderson et al. 2012b; Khramtsova et al. 2013; Anderson et al. 2014; Makai et al. 2017; Topchieva et al. 2017) is a powerful tool to advance further interpretation of observational data and to relate them to results of numerical investigations. This is important as there are still some key questions, which lack definite answers. We mention briefly some of them.

It is still not clear how the object's size is related to its age. Three varieties of HII regions are distinguished, namely:

- (1) ultracompact and hypercompact HII regions (size less than 0.1 pc, electron density $> 10^4 \text{ cm}^{-3}$);
- (2) classic HII regions (size of the order of a few parsecs, electron density $\sim 10^2 \text{ cm}^{-3}$);
- (3) giant HII regions (size of the order of 100 pc, density $< 30 \text{ cm}^{-3}$).

It is possible that they all represent different stages of a single process. Specifically, it has been suggested in Zinnecker & Yorke (2007) that smaller and denser HII regions are young, while less dense and more extended

HII regions are older. However, the exact evolutionary relationships between HII regions of various kinds are still unclear, and we cannot be certain if they exist at all.

The second problem, which is hard to solve, is the identification of a star that ionizes a given object. A standard suggestion, which is routinely adopted in various models, is the central location of the star (Gail & Sedlmayr 1979; Arthur et al. 2004; Draine 2011a; Pavlyuchenkov et al. 2013; Akimkin et al. 2015, 2017). However, the ionizing star in an HII region may reside not only in its center, but also on the periphery and even beyond the object. The latter two morphologies are usually referred to as champaign flow and blister, respectively. Examples of such morphologies were found in Sh2-212 by Deharveng & Zavagno (2008) and Deharveng et al. (2008), in the Orion Nebula by O’Dell & Yusef-Zadeh (2000), and in several bipolar HII regions by Deharveng et al. (2012) and Deharveng et al. (2015). It is also possible that some HII regions are excited by several OB stars, e.g. RCW 79 (Martins et al. 2010).

To attack all these problems we need a self-consistent evolutionary model of HII regions, which is able to reproduce distributions of density, temperature, velocity and molecular abundances simultaneously. On the other hand, we need thoroughly analyzed observational data, suitable for comparison with theoretical results. In this paper we present an analysis of photometry of HII regions in order to construct a large sample of objects, which can be used for comparison both with results of numerical simulations and with results from other studies of HII regions and complexes (Anderson et al. 2012b; Khramtsova et al. 2013).

As an example of application of the presented photometric catalog, we estimate the polycyclic aromatic hydrocarbon (PAH) mass fraction (q_{PAH}) and intensity of ultraviolet (UV) irradiation in the studied objects, using the grid of models by Draine & Li (2007). Also we analyze possible differences in flux ratios between Galactic HII regions with resolved structure and extragalactic HII complexes, which are spatially unresolved.

In Section 2 we describe data processing. Section 3 contains photometric analysis of IR ring nebula images. In Section 4 results are presented and discussed.

2 DATA PROCESSING

We use a catalog presented in the work of Topchieva et al. (2017). The New GPS 20 cm survey, created using the MAGPIS database of radio images of regions with Galactic coordinates $|b_{\text{gal}}| < 0.8^\circ$, $5^\circ < l_{\text{gal}} < 48.5^\circ$, was used as the basis for this study. We identified compact sources of radio emission among the ob-

jects in this survey, toward which we performed a visual search of objects, which look like rings at $8\ \mu\text{m}$ and contain IR emission at $24\ \mu\text{m}$ and radio emission at 20 cm in their interiors. The catalog is based on IR survey data of 8 and $24\ \mu\text{m}$, obtained with the IRAC (Fazio et al. 2004) and MIPS (Rieke et al. 2004) instruments on the *Spitzer Space Telescope*. At longer wavelengths we used data from the *Herschel Space Observatory* science archive. Images in 70 and $160\ \mu\text{m}$ were acquired with the PACS instrument (Poglitsch et al. 2010), while images in $250\ \mu\text{m}$, $350\ \mu\text{m}$ and $500\ \mu\text{m}$ were recorded with the SPIRE instrument (Griffin et al. 2010). The total number of selected objects is 99. Four of them have not been identified in previous surveys of IR ring nebulae. Further, they are designated as TWKK. Unlike other catalogs, this catalog is specifically prepared as a resource for comparison with results of 1D spherically symmetric hydrodynamical computations. Thus, it only includes bright sources with a more or less regular structure.

Qualitatively, we can assume that at $8\ \mu\text{m}$, the major contribution to the object emission comes from IR bands attributed to PAHs. At $24\ \mu\text{m}$ the main sources of emission are presumable stochastically heated very small grains along with, probably, hot large grains (Paladini et al. 2012). At longer wavelengths emission is mostly generated by colder large grains (Draine 2011b). At $250\ \mu\text{m}$, $350\ \mu\text{m}$ and $500\ \mu\text{m}$ due to low angular resolution it is hard to distinguish between the object emission and the background (and foreground) emission (Anderson et al. 2012a), but we still consider these bands in our study, trying to make the best effort in removing the background contribution.

2.1 Aperture Photometry

Before the flux estimation, photometry data in all wavebands have been convolved to the same resolution using kernels from Aniano et al. (2011). We keep the original pixel size and take it into account, when computing fluxes.

Size and location of a source aperture were selected using $8\ \mu\text{m}$ data based on the catalog from Topchieva et al. (2017). In order to estimate emission fluxes we need to get rid of background radiation and radiation from other sources, which are not related to the studied object (for example, stars and Galactic background radiation). This is why we clean the image with respect to point sources and subtract the background, estimated using a separate aperture, which is located in the darkest spot of the map beyond the object aperture. We believe that the background value measured at a brighter location or a




Fig. 1 A map showing locations of the source and background apertures at $8\ \mu\text{m}$ for the object CN67. A *white circle* shows the source aperture, while a *red circle* shows the aperture used for background estimation.

background value, averaged over the entire image, can significantly affect the estimate of a useful source signal. Background aperture size depends on the extent of the area selected for background estimation (Fig. 1).

In all wavebands other than $8\ \mu\text{m}$ (24, 70, 160, 250, 350 and $500\ \mu\text{m}$) for both source flux estimation and background estimation, we use the same apertures as at $8\ \mu\text{m}$. Custom-made Python scripts are utilized to compute total source fluxes in the above bands.

Apart from the HII region fluxes themselves, we also consider flux ratios, or color indices, $[F_{24}/F_8]$, $[F_{70}/F_{24}]$, $[F_{160}/F_{24}]$ and $[F_{160}/F_{70}]$, and compare them to the criteria suggested by Anderson et al. (2012a). The authors have shown that these ratios can be used to discriminate between unresolved planetary nebulae and HII regions. Below we check whether the criteria developed by Anderson et al. (2012a) can be applied to HII regions with spatially resolved structure. Also we relate our results in terms of flux ratios to those from the work of Khramtsova et al. (2013), where unresolved extragalactic HII regions have been considered.

2.2 PAH Mass Fraction and UV Field Intensity

Apart from simply computing flux ratios, we performed a somewhat more sophisticated analysis and estimated a PAH mass fraction, q_{PAH} , using a grid of models from Draine & Li (2007). The value of q_{PAH} is an important parameter in at least two respects. First, PAH emission is often considered as an indicator of the star formation rate, because corresponding transitions are excited by UV photons, which presumably trace the number of young stars. However, the relation between strength of

the IR bands and the intensity of UV radiation can be non-trivial as UV photons both excite PAHs and destroy them. Thus, it is important to consider how q_{PAH} behaves on various spatial scales. Specifically, a small value of q_{PAH} may indicate that organic dust particles are destroyed in HII regions (Madden et al. 2006; Leboutteiller et al. 2007). Second, the evolution of PAHs in star-forming regions is important in the context of a general evolution of organic matter in the universe.

The grid of models from Draine & Li (2007) can be used to determine a UV field intensity U in these regions. In this formalism, the radiation fields of the region under consideration are assumed to consist of two components, the minimum radiation field with intensity U_{min} and the enhanced radiation field with intensity U , distributed between U_{min} and some maximum value, U_{max} . We consider a simplified situation of an isolated HII region, so we assume that there is a single value of radiation intensity, U_{min} (in other words, $U_{\text{max}} = U_{\text{min}}$), and denote it simply as U . This is also an important parameter as UV photons are crucial for thermal balance and dynamical evolution both inside the HII region and at its border, where a so-called photon-dominated region (PDR) is located. Both parameters are related to each other as UV photons both destroy PAH particles and excite them, so that they can generate IR emission bands. Also, in PDRs 0.1%–1% of absorbed UV photons are transferred to suprathermal ($\sim 1\ \text{eV}$) photoelectrons, being ejected from dust and PAH particles. These electrons heat the gas, so an estimate of q_{PAH} is needed to evaluate the contribution of PAHs to the thermal balance in these regions. In the grid of models by Draine & Li (2007), UV field intensity is measured in units of the intensity of the average radiation field in the solar vicinity.

3 RESULTS

Performed flux measurements, presented in Table 1, were used to construct spectral energy distributions (SEDs) for all the studied objects. We do see some variations in the SED shapes for different regions. Most objects have SEDs with the expected outline, that is, maximum emission at 70 and $160\ \mu\text{m}$ and shallow decrease toward $500\ \mu\text{m}$, but there are some noticeable exceptions (Fig. 2). For example, objects TWKK2 and MWP1G034088+004405 show “flat” profiles, indicative of the excess presence of warmer dust. In objects MWP1G024019+001902 and S15, radiation flux decreases toward $250\ \mu\text{m}$, which could mean incorrect background subtraction for these objects. We hope that HII region modeling will help to clarify these issues.




Fig. 2 SEDs for objects TWKK2, MWP1G034088+004405, MWP1G024019+001902 and S15. *Lines* show the SED without background subtraction, while *triangles* show flux values with background taken into account.

The second column of Table 1 contains the object designation. Galactic coordinates of the object are shown in columns 3 and 4. Fluxes in MJy are given in columns 5–11. Finally, in columns 12 and 13 we show estimates for PAH mass fraction and UV field intensity.

Analysis of the PAH mass fraction shows that almost all the objects, except for N14, N51 and MWP1G03080+001100S, are characterized by q_{PAH} less than 0.47%. This is an upper limit as the grid of models by Draine & Li (2007) does not contain data for smaller values of q_{PAH} . Nevertheless, this result agrees with our expectation that q_{PAH} in HII regions is smaller than the average value for the Milky Way galaxy (that is, about a few percent) due to PAH destruction in HII regions (Madden et al. 2006; Lebouteiller et al. 2007). Note that there are some objects with q_{PAH} value of about 4% or greater. A case by case examination shows that these are regions with more complex morphology and with a significant unaccounted background contribution.

Values of UV field intensity U are surprisingly low, from 1 to 10, which is lower than would be expected for the immediate vicinity of a massive star. However, this value is mostly determined by the far-IR part of the spec-

trum. In our objects this emission comes from the outer rings, which are located far away from the ionizing stars. A more detailed picture will arise when we will analyze SED radial variations. Also, it would be interesting to relate the U value to the linear size of the object, checking a naive assumption that the object size can serve as a qualitative measure of its age. However, distances ought to be known for this, and kinematic distance estimates are only available for 11 objects from our sample.

It would be interesting to relate obtained values of U and q_{PAH} to each other. Specifically, it has been shown in the work of Khramtsova et al. (2013) that the PAH mass fraction tends to be smaller in star-forming complexes with greater U_{min} . Obviously, we do not see a similar trend in our data. There are at least two reasons for this. First, we would like to emphasize that in most objects we were only able to get an upper limit for q_{PAH} . Thus, strictly speaking, we cannot say for certain whether or not q_{PAH} (anti)correlates with U . It has been argued in Khramtsova et al. (2013) that the ratio of fluxes at 8 and 24 μm can be used as a substitute for q_{PAH} , but this is only true when the fluxes are computed for large star-forming complexes. In the presented study this does not

Table 1 HII Region Parameters

No.	Object	$l_{\text{gal}} (^{\circ})$	$b_{\text{gal}} (^{\circ})$	F_8 (MJy)	F_{24} (MJy)	F_{70} (MJy)	F_{160} (MJy)	F_{250} (MJy)	F_{350} (MJy)	F_{500} (MJy)	q_{PAH}	U
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
1	S15 ³	343.916	-0.648	6.28(-5)	9.55(-4)	4.45(-2)	5.09(-2)	7.43(-4)	1.21(-3)	5.06(-4)	0.47	10.0
2	S21 ³	341.358	-0.288	1.54(-5)	9.45(-6)	2.19(-3)	3.60(-3)	2.39(-4)	1.06(-4)	3.88(-5)	0.47	4.0
3	S44 ³	334.524	0.820	1.17(-4)	1.99(-4)	5.36(-2)	6.01(-2)	1.48(-3)	5.76(-4)	1.86(-4)	0.47	10.0
4	S123 ³	312.978	-0.433	6.74(-5)	6.56(-5)	3.30(-2)	6.46(-2)	1.09(-3)	4.60(-4)	1.65(-4)	0.47	5.0
5	S145 ³	308.717	0.623	7.86(-4)	2.47(-3)	5.44(-1)	4.93(-1)	6.85(-3)	4.47(-3)	3.01(-3)	0.47	12.0
6	S167 ³	301.627	-0.345	2.65(-5)	1.17(-4)	9.15(-3)	2.98(-2)	2.82(-3)	2.60(-3)	2.30(-3)	0.47	1.5
7	CN67 ³	5.526	0.037	1.18(-5)	1.65(-5)	7.88(-3)	1.35(-2)	2.33(-4)	9.81(-5)	3.52(-5)	0.47	4.0
8	CN77 ³	6.139	-0.640	1.15(-5)	1.94(-4)	6.47(-3)	5.48(-3)	7.48(-4)	2.82(-4)	9.33(-5)	0.47	20.0
9	CN79 ³	6.202	-0.334	2.94(-5)	5.89(-5)	9.41(-3)	1.40(-2)	9.77(-4)	4.49(-4)	1.60(-4)	0.47	7.0
10	CN111 ³	8.311	-0.086	2.88(-5)	5.12(-5)	2.51(-2)	4.85(-2)	8.57(-4)	3.90(-4)	1.47(-4)	0.47	3.0
11	MWP1G008430-002800S ²	8.431	-0.276	7.82(-6)	1.09(-5)	8.21(-3)	9.48(-3)	2.94(-4)	1.39(-4)	5.14(-5)	0.47	10.0
12	CN116 ³	8.476	-0.277	7.63(-6)	9.18(-6)	6.70(-3)	7.88(-3)	1.33(-4)	5.83(-5)	2.14(-5)	0.47	10.0
13	N4 ³	11.893	0.747	2.28(-4)	2.47(-4)	8.18(-2)	8.82(-2)	1.25(-3)	9.02(-4)	6.43(-4)	0.47	10.0
14	MWP1G012590-000900S ²	12.595	-0.090	2.26(-6)	3.78(-6)	3.54(-3)	4.33(-3)	6.43(-5)	2.50(-5)	8.20(-6)	0.47	10.0
15	MWP1G012630-000100S ²	12.633	-0.017	7.63(-6)	1.15(-5)	1.23(-2)	2.18(-2)	4.59(-4)	2.24(-4)	8.45(-5)	0.47	3.0
16	N8 ³	12.805	-0.312	2.43(-6)	3.35(-6)	1.19(-3)	2.12(-3)	1.71(-4)	8.00(-5)	3.97(-5)	0.47	3.0
17	MWP1G013213-001410 ²	13.213	-0.141	2.42(-5)	4.90(-5)	4.03(-2)	5.82(-2)	1.14(-3)	5.01(-4)	1.74(-4)	0.47	5.0
18	N13 ³	13.899	-0.014	5.79(-6)	9.29(-6)	1.59(-3)	2.49(-3)	1.69(-4)	7.49(-5)	2.81(-5)	0.47	5.0
19	N14 ³	14.000	-0.136	4.67(-4)	4.25(-4)	4.55(-2)	4.64(-2)	2.65(-3)	1.08(-3)	3.76(-4)	3.9–4.6	15.0
20	G014.175+0.024 ^{6,1}	14.175	0.022	2.06(-6)	4.72(-6)	2.81(-3)	3.53(-3)	5.56(-5)	2.27(-5)	7.51(-6)	0.47	10.0
21	MWP1G014210-001100S ²	14.206	-0.110	2.03(-6)	4.33(-6)	1.70(-3)	1.37(-3)	3.90(-5)	1.57(-5)	5.20(-6)	0.47	25.0
22	MWP1G014390-000200S ²	14.388	-0.024	9.87(-6)	8.41(-6)	4.68(-3)	2.81(-3)	1.09(-4)	4.31(-5)	1.46(-5)	0.47	25.0
23	MWP1G014480-000000S ²	14.490	0.022	4.85(-6)	7.33(-6)	6.54(-3)	1.15(-2)	2.20(-4)	1.01(-4)	3.62(-5)	0.47	4.0
24	MWP1G016390-001400S ²	16.391	-0.138	1.50(-6)	2.63(-6)	1.93(-3)	2.43(-3)	3.96(-5)	1.63(-5)	5.53(-6)	0.47	8.0
25	MWP1G016429-001984 ²	16.431	-0.201	2.11(-5)	5.59(-5)	2.94(-2)	4.50(-2)	7.66(-4)	3.30(-4)	1.14(-4)	0.47	5.0
26	MWP1G016560+000056 ²	16.560	0.002	2.11(-6)	1.59(-6)	1.83(-3)	3.96(-3)	6.95(-5)	2.96(-5)	1.05(-5)	0.47	2.5
27	MWP1G017626+000493 ²	17.625	0.048	4.55(-6)	3.22(-6)	2.32(-3)	5.56(-3)	1.03(-4)	5.14(-5)	2.14(-5)	0.47	2.5
28	TWKK1 ⁴	17.805	0.074	1.80(-6)	9.74(-7)	6.62(-4)	9.98(-4)	1.24(-5)	8.90(-6)	3.27(-6)	0.47	4.0
29	N20 ³	17.918	-0.687	7.81(-7)	1.38(-5)	5.94(-4)	9.54(-4)	2.17(-4)	1.06(-4)	3.85(-5)	0.47	4.0
30	MWP1G018440+000100S ²	18.442	0.013	6.87(-7)	1.68(-6)	1.43(-3)	1.68(-3)	3.23(-5)	1.67(-5)	6.68(-6)	0.47	10.0
31	MWP1G018580+003400S ²	18.582	0.345	2.24(-6)	7.81(-6)	1.66(-4)	1.95(-4)	1.42(-4)	6.68(-5)	2.57(-5)	4.60	12.0–15.0
32	N23 ³	18.679	-0.237	9.81(-6)	1.98(-5)	9.11(-3)	9.03(-3)	2.95(-4)	1.42(-4)	5.46(-5)	0.47	12.0
33	MWP1G018743+002521 ²	18.748	0.256	1.14(-5)	2.52(-5)	9.40(-3)	1.30(-2)	3.26(-5)	9.15(-5)	3.37(-5)	0.47	5.0–7.0
34	MWP1G020387-000156 ²	20.388	-0.017	6.80(-6)	4.68(-6)	4.76(-3)	7.25(-3)	9.21(-5)	8.67(-5)	3.71(-5)	0.47	5.0
35	MWP1G02100-000500S ²	21.005	-0.054	1.53(-6)	3.45(-6)	2.51(-4)	4.13(-4)	7.00(-6)	2.98(-6)	1.08(-6)	0.47	3.0
36	N28 ³	21.351	-0.137	2.56(-5)	2.46(-5)	3.46(-3)	3.76(-3)	2.18(-4)	1.88(-4)	7.00(-5)	0.47	10.0
37	N31 ³	23.842	0.098	5.24(-6)	7.14(-6)	6.40(-3)	8.22(-3)	1.33(-4)	5.80(-5)	2.10(-5)	0.47	7.0
38	MWP1G023849-001251 ²	23.848	-0.127	7.82(-6)	1.01(-5)	5.51(-3)	7.15(-3)	1.07(-4)	4.65(-5)	1.69(-5)	0.47	7.0
39	MWP1G023881-003497 ²	23.881	-0.350	4.37(-6)	8.28(-6)	5.50(-4)	5.14(-4)	4.68(-6)	7.61(-6)	2.61(-6)	0.47–1.2	12.0
40	N32 ³	23.904	0.070	5.70(-6)	1.35(-5)	1.16(-2)	1.56(-2)	2.65(-4)	1.24(-4)	4.78(-5)	0.47	5.0
41	MWP1G023982-001096 ²	23.982	-0.110	1.53(-6)	3.22(-6)	2.40(-4)	1.58(-4)	1.11(-5)	4.97(-6)	1.79(-6)	2.5–4.6	25.0
42	MWP1G024019+001902 ²	24.043	0.204	7.72(-7)	1.69(-6)	1.28(-4)	1.22(-4)	9.80(-7)	3.56(-6)	1.36(-6)	0.47	15.0
43	MWP1G024149-000060 ²	24.153	-0.011	3.12(-6)	3.94(-6)	3.15(-3)	3.27(-3)	1.59(-4)	3.30(-5)	1.22(-5)	0.47	12.0–15.0
44	N33 ³	24.215	-0.044	5.68(-6)	1.83(-5)	8.11(-3)	8.44(-3)	1.23(-4)	5.39(-5)	1.93(-5)	0.47	12.0–15.0
45	TWKK3 ⁴	24.424	0.220	1.76(-5)	3.86(-5)	1.70(-2)	2.37(-2)	5.00(-4)	2.47(-4)	9.05(-5)	0.47	7.0
46	TWKK2 ⁴	24.460	0.506	3.97(-7)	7.39(-6)	3.38(-4)	3.46(-4)	4.99(-5)	1.95(-5)	6.38(-6)	0.47	20.0
47	MWP1G024500-002400 ²	24.502	-0.237	1.24(-5)	2.00(-5)	7.66(-3)	8.62(-3)	1.62(-4)	6.69(-5)	6.15(-6)	0.47	15.0
48	MWP1G024558-001329 ²	24.558	-0.133	3.25(-5)	2.09(-5)	3.39(-3)	5.52(-3)	4.02(-4)	1.92(-4)	7.16(-5)	1.12	5.0
49	MWP1G024649-001131 ²	24.651	-0.078	2.93(-6)	2.81(-6)	2.08(-3)	1.60(-3)	1.78(-5)	7.85(-6)	3.12(-6)	0.47	20.0
50	MWP1G01024699-001486 ²	24.700	-0.148	1.56(-5)	2.66(-5)	1.14(-2)	9.54(-3)	1.13(-4)	3.37(-5)	1.03(-5)	0.47	25.0
51	MWP1G024731+001580 ²	24.736	0.158	1.63(-5)	2.23(-5)	1.07(-2)	1.27(-2)	3.28(-4)	1.40(-4)	4.82(-5)	0.47	10.0
52	MWP1G024920+000800 ²	24.922	0.078	3.32(-6)	8.33(-6)	7.03(-3)	1.13(-2)	2.10(-4)	9.74(-5)	3.50(-5)	0.47	5.0
53	MWP1G025155+000609 ²	25.155	0.061	2.97(-5)	2.72(-5)	1.12(-2)	1.02(-2)	1.13(-4)	7.42(-5)	2.56(-5)	0.47	15.0
54	MWP1G025723+00058 ²	25.724	0.058	1.16(-5)	1.78(-5)	9.62(-3)	9.15(-3)	1.15(-4)	4.41(-5)	1.48(-5)	0.47	15.0
55	MWP1G025730-000200S ²	25.726	-0.027	3.02(-6)	2.88(-6)	1.97(-3)	6.83(-4)	4.05(-5)	1.54(-5)	4.39(-6)	0.47	—
56	N42 ³	26.329	-0.071	1.46(-5)	1.54(-5)	1.11(-2)	1.57(-2)	2.46(-4)	1.04(-4)	3.58(-5)	0.47	5.0–7.0
57	N43 ³	26.595	0.095	1.39(-5)	2.06(-5)	1.09(-2)	1.44(-2)	2.37(-4)	1.04(-4)	3.73(-5)	0.47	7.0
58	MWP1G026720+001700S ²	26.722	0.173	4.88(-6)	5.00(-6)	3.51(-3)	4.13(-3)	4.12(-5)	2.97(-5)	1.03(-5)	0.47	10.0
59	G027.492+0.192 ⁶	27.496	0.197	3.15(-5)	8.70(-5)	3.84(-2)	3.20(-2)	3.94(-4)	1.51(-4)	5.14(-5)	0.47	20.0
60	MWP1G02671+00300S ²	27.613	0.028	2.65(-6)	2.28(-6)	1.47(-3)	1.40(-3)	3.62(-5)	1.47(-5)	5.04(-6)	0.47–1.2	12.0–15.0
61	MWP1G027905-000079 ²	27.904	-0.009	9.35(-6)	1.11(-5)	7.77(-3)	9.53(-3)	1.45(-4)	5.78(-5)	1.92(-5)	0.47	7.0–10.0
62	G027.9334+00.2056 ^{6,5}	27.931	0.205	3.11(-6)	9.73(-6)	6.03(-3)	6.81(-3)	9.49(-5)	3.76(-5)	1.20(-5)	0.47	10.0
63	MWP1G027981+000753 ²	27.981	0.073	1.96(-5)	2.50(-5)	4.36(-3)	4.41(-3)	2.41(-4)	1.67(-4)	5.74(-5)	0.47	10.0–12.0
64	MWP1G028160-000300S ²	28.160	-0.046	8.14(-6)	4.16(-6)	4.97(-3)	7.72(-3)	1.14(-4)	4.28(-5)	1.38(-5)	0.47	5.0
65	N49 ³	28.827	-0.229	9.90(-5)	1.86(-4)	1.94(-2)	3.42(-2)	1.89(-3)	9.77(-4)	3.46(-4)	0.47	3.0–4.0
66	MWP1G029136-001438 ²	29.134	-0.144	3.43(-6)	4.20(-6)	3.07(-3)	3.60(-3)	5.50(-5)	2.38(-5)	8.68(-6)	0.47	10.0

Table 1 — Continued.

No.	Object	$l_{\text{gal}} (^{\circ})$	$b_{\text{gal}} (^{\circ})$	F_8 (MJy)	F_{24} (MJy)	F_{70} (MJy)	F_{160} (MJy)	F_{250} (MJy)	F_{350} (MJy)	F_{500} (MJy)	q_{PAH}	U
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
67	N51 ³	29.156	-0.259	5.12(-5)	1.05(-4)	4.80(-3)	5.64(-3)	2.54(-4)	1.28(-4)	6.91(-5)	3.9–4.6	12.0
68	MWP1G030020–000400S ²	30.022	-0.041	2.95(-5)	4.65(-5)	8.18(-3)	8.26(-3)	4.49(-4)	1.71(-4)	5.35(-5)	0.47	15.0
69	MWP1G030250+002413 ²	30.251	0.240	1.31(-5)	1.29(-5)	1.30(-2)	1.48(-2)	1.82(-4)	7.61(-5)	2.67(-5)	0.47	10.0
70	MWP1G03080+001100S ²	30.378	0.111	7.89(-6)	1.25(-5)	7.89(-3)	8.59(-3)	1.25(-4)	5.08(-5)	1.76(-5)	0.47	10.0
71	MWP1G030381–001074 ²	30.381	-0.109	3.39(-5)	4.56(-5)	2.71(-2)	2.05(-2)	2.25(-4)	1.61(-4)	6.58(-5)	0.47	20.0
72	MWP1G031066+000485 ²	31.071	0.049	4.63(-6)	1.01(-5)	6.27(-3)	3.76(-3)	3.36(-5)	1.83(-5)	2.27(-6)	0.47	25.0
73	MWP1G032057+000783 ²	32.055	0.076	1.76(-5)	2.97(-5)	3.25(-2)	4.20(-2)	7.57(-4)	3.63(-4)	1.32(-4)	0.47	7.0–10.0
74	N55 ³	32.101	0.091	3.78(-5)	3.30(-5)	7.68(-3)	1.05(-2)	1.65(-4)	7.03(-5)	2.46(-5)	0.47	3.0–5.0
75	MWP1G032731+002120 ²	32.730	0.212	7.42(-6)	7.64(-6)	4.72(-3)	9.30(-3)	1.60(-4)	7.30(-5)	2.76(-5)	0.47	3.0–4.0
76	N57 ³	32.761	-0.149	3.41(-6)	2.25(-6)	2.14(-3)	3.73(-3)	5.51(-5)	2.19(-5)	7.68(-6)	0.47	3.0–4.0
77	N60 ³	33.815	-0.149	9.26(-6)	1.14(-5)	7.95(-3)	8.24(-3)	1.13(-4)	5.26(-5)	3.45(-5)	0.47	10.0
78	MWP1G034088+004405 ²	34.087	0.441	1.31(-6)	2.43(-5)	9.58(-4)	9.94(-4)	1.55(-4)	6.15(-5)	2.14(-5)	0.47	10.0–12.0
79	MWP1G034680+000600S ²	34.684	0.067	2.42(-6)	4.31(-6)	3.04(-3)	3.26(-3)	4.92(-5)	1.99(-5)	6.65(-6)	0.47	10.0–12.0
80	N67 ³	35.544	0.012	1.07(-5)	2.80(-5)	9.25(-3)	6.60(-3)	2.28(-4)	8.54(-5)	2.98(-5)	0.47	20.0
81	MWP1G037196–004296 ²	37.195	-0.429	6.24(-6)	1.15(-5)	1.50(-3)	1.59(-3)	2.28(-5)	2.17(-5)	8.09(-6)	0.47	10.0–12.0
82	MWP1G037261–000809 ²	37.258	-0.078	1.96(-5)	2.63(-5)	1.24(-2)	1.45(-2)	1.74(-4)	5.65(-5)	2.55(-5)	0.47	10.0
83	MWP1G037349+006876 ²	37.351	0.688	1.10(-5)	2.11(-4)	8.58(-3)	9.27(-3)	1.27(-4)	5.17(-5)	1.93(-5)	0.47	10.0
84	N70 ³	37.750	-0.113	5.75(-6)	2.46(-5)	1.05(-2)	1.21(-2)	1.73(-4)	7.28(-5)	2.52(-5)	0.47	10.0
85	G038.550+1648 ⁶	38.551	0.162	2.69(-6)	9.07(-6)	3.84(-3)	4.05(-3)	6.18(-5)	2.43(-5)	8.43(-6)	0.47	10.0–12.0
86	N73 ³	38.736	-0.140	2.10(-5)	1.31(-5)	3.68(-3)	7.53(-3)	5.28(-4)	2.23(-4)	7.92(-5)	0.47	3.0–4.0
87	N78 ³	41.228	0.169	1.92(-6)	1.78(-6)	1.13(-3)	1.70(-3)	2.67(-5)	1.11(-5)	4.27(-6)	0.47	5.0–7.0
88	G041.378+0.035 ^{6,1}	41.378	0.034	6.48(-6)	1.46(-5)	6.13(-3)	5.97(-3)	9.20(-5)	3.81(-5)	1.31(-5)	0.47	15.0–20.0
89	N79 ³	41.513	0.031	2.01(-5)	4.46(-5)	2.22(-2)	2.32(-2)	3.12(-4)	1.26(-4)	4.33(-5)	0.47	10.0–12.0
90	TWKK4 ⁴	41.595	0.160	1.42(-6)	1.89(-6)	9.89(-4)	1.10(-3)	1.58(-5)	6.42(-6)	2.18(-6)	0.47	10.0–12.0
91	N80 ³	41.932	0.033	1.42(-5)	1.55(-5)	2.12(-3)	3.62(-3)	2.64(-4)	1.31(-4)	5.57(-5)	0.47	3.0–4.0
92	N89 ³	43.739	0.114	5.14(-6)	8.00(-5)	3.07(-3)	3.19(-3)	4.45(-5)	2.72(-5)	1.70(-5)	0.47	10.0–12.0
93	N90 ³	43.774	0.060	2.39(-5)	2.05(-5)	7.07(-4)	2.30(-3)	3.15(-4)	1.06(-4)	4.39(-5)	0.47	1.0–1.5
94	MWP1G045540+000000S ²	45.544	-0.005	5.72(-6)	1.33(-5)	5.87(-3)	4.48(-3)	5.86(-5)	2.35(-5)	7.87(-6)	0.47	20.0–25.0
95	N96 ³	46.949	0.371	2.96(-6)	6.78(-6)	6.97(-4)	7.35(-4)	3.89(-5)	2.77(-5)	1.13(-5)	0.47	10.0
96	N92 ³	47.027	0.218	3.21(-5)	3.16(-5)	4.58(-3)	9.43(-3)	7.49(-4)	3.54(-4)	1.33(-4)	0.47–1.2	3.0–4.0
97	MWP1G048422+001173 ²	48.422	0.116	9.92(-6)	7.56(-6)	5.56(-3)	8.46(-3)	1.14(-4)	4.37(-5)	1.48(-5)	0.47	3.0–5.0
98	N102 ³	49.697	-0.164	1.14(-5)	6.40(-5)	2.05(-2)	9.30(-3)	1.33(-4)	6.45(-5)	6.49(-5)	0.47	25.0
99	N121 ³	55.444	0.887	1.65(-6)	1.05(-5)	3.17(-4)	1.10(-3)	2.43(-5)	1.92(-5)	1.54(-5)	0.47	1.0–1.5

Notes: The numbers between parentheses, $a(b)$, mean $a \times 10^b$. Objects are taken from ¹Becker et al. (1994), ²Simpson et al. (2012), ³Churchwell et al. (2006), ⁴new objects, ⁵Urquhart et al. (2009) and ⁶Egan et al. (2003).

work as in our objects 8 μm emission and 24 μm emission are spatially separated from each other. Thus, we do not see any correlation between U and F_8/F_{24} either. Second, we interpret small upper limits for q_{PAH} as a signature of PAH destruction in the vicinity of an ionizing star or stars, so we may expect a clearer relation between q_{PAH} and U closer to the source of UV radiation. On the other hand, the derived U value is to a large degree determined by far IR emission, which mostly comes from the region periphery. This is why our derived U values are close to those presented in Khramtsova et al. (2013), even though our U is not an exact equivalent of U_{min} from Khramtsova et al. (2013). The latter value is more representative of the space between individual HII regions in a large star-forming complex.

In general, a major difference between the work of Khramtsova et al. (2013) and the present study is in vastly different spatial scales. In most regions presented in Khramtsova et al. (2013) q_{PAH} is about a few percent, which is significantly higher than in our regions. This implies that PAH destruction proceeds locally, in individual HII regions, while most emission at 8 μm comes from

more extended inter-region areas, which are not probed in the present study.

In Figure 3 we relate our flux ratios [F_{24}/F_8], [F_{70}/F_{24}], [F_{160}/F_{24}] and [F_{160}/F_{70}] to the criteria suggested in Anderson et al. (2012a) and similar ratios for extragalactic HII complexes presented by Khramtsova et al. (2013) (square brackets indicate the logarithm of the corresponding ratio). In the top right corner of each panel we indicate the value of the flux ratio logarithm, which was indicated in Anderson et al. (2012a) as discriminating between HII regions and planetary nebulae. These values are also marked with vertical lines in each panel.

Obviously, most of our objects (hashed bars) and extragalactic HII regions (empty bars) satisfy the Anderson et al. (2012a) criteria and are indeed HII regions. However, there are few objects which are definitely HII regions, but would have been classified as planetary nebulae by Anderson et al. (2012a) constraints. Specifically, in some objects, values of [F_{24}/F_8] ratios are greater than what are expected in HII regions, while for some objects the values of [F_{70}/F_{24}], [F_{160}/F_{24}] and [F_{160}/F_{70}] ratios




Fig. 3 Flux ratios for 99 HII regions studied in this paper (*hashed bars*) and for extragalactic HII complexes studied in Khramtsova et al. (2013) (*empty bars*). A *black vertical line* indicates a value discriminating between HII regions and planetary nebulae according to Anderson et al. (2012a). The value is also shown in the *top right corner* of each panel. a) $[F_{24}/F_8]$; b) $[F_{70}/F_{24}]$; c) $[F_{160}/F_{24}]$; d) $[F_{160}/F_{70}]$. All the fluxes from this work are used with background subtraction.

are smaller than values expected in HII regions. This emphasizes that simple photometric criteria may fail, when they are applied to spatially resolved objects. We believe that the reason is two-fold. First, we draw the outer boundary of an object by relying on the location of the outer 8 μm ring. However, 8 μm emission allows drawing a boundary of an HII region only in the sense that it traces the location of a dense shell swept up by ionization and shock fronts. At the same time, in nearly all cases we also see a somewhat fainter 8 μm emission, which extends well beyond the dense shell but is also related to the considered object. In other words, when we compare wide-scale maps of emission at 8 and 24 μm , we see that 8 μm emission is more extended than 24 μm emission. Thus, we may inadvertently miss some 8 μm emission which is located beyond the ring but still belongs to the object, while 24 μm emission is accounted for entirely. On the other hand, due to lower angular resolution at longer wavelengths, some emission at these wavelengths may leak out of the 8 μm -based aperture.

As for fluxes, computed by Khramtsova et al. (2013) for extragalactic sources, their differences both from our data and from the Anderson et al. (2012a) criteria

are more significant, especially for the $[F_{70}/F_{24}]$ and $[F_{160}/F_{24}]$ ratios. This is probably again related to a drastically different spatial scale probed in the work of Khramtsova et al. (2013). Star-forming complexes studied in that paper have typical linear sizes of a few hundred pc. In this case a single aperture includes both numerous individual HII regions and material between them. As we have mentioned above and as had been found earlier in Bendo et al. (2008), emission at 24 μm is more compact than emission at other wavelengths, while emission at the far-IR range is more diffuse. Thus, when an aperture corresponds to a large linear scale, we may expect a more significant contribution from emission at 70 and 160 μm . This is why the $[F_{70}/F_{24}]$ and $[F_{160}/F_{24}]$ flux ratios are greater in the work of Khramtsova et al. (2013) than in the present paper.

4 CONCLUSIONS

The following results are presented in this work;

- (1) Total fluxes at 8, 24, 70, 160, 250, 350 and 500 μm are estimated for 99 HII regions. This information

can later be used for comparison with results of theoretical computations.

- (2) A PAH mass fraction q_{PAH} is estimated for these regions. In most regions we have only been able to obtain upper limits for q_{PAH} , showing that the actual values are smaller than 0.47%. This value is much lower than the average Galactic PAH mass fraction, which is about a few percent. We argue that this is a signature of local PAH destruction in HII regions.
- (3) Flux ratios $[F_{24}/F_8]$, $[F_{70}/F_{24}]$, $[F_{160}/F_{24}]$ and $[F_{160}/F_{70}]$ are estimated. It is shown that in some cases the criteria, suggested in Anderson et al. (2012a) to distinguish between HII regions and planetary nebulae, may fail when applied to spatially resolved objects.
- (4) Systemic differences with flux measurements in extragalactic HII complexes (Khrantsova et al. 2013) are caused by significantly different spatial scales (Samus & Li 2018).

Acknowledgements This study is supported by Program 7 of the Presidium of the RAS, “Transitional and Explosive Processes in Astrophysics,” and the RFBR (Grant 17-02-00521). The Astropy (Astropy Collaboration et al. 2013) package has been used to obtain the presented results.

References

- Akimkin, V. V., Kirsanova, M. S., Pavlyuchenkov, Y. N., & Wiebe, D. S. 2015, *MNRAS*, 449, 440
- Akimkin, V. V., Kirsanova, M. S., Pavlyuchenkov, Y. N., & Wiebe, D. S. 2017, *MNRAS*, 469, 630
- Anderson, L. D., Bania, T. M., Balsler, D. S., et al. 2014, *ApJS*, 212, 1
- Anderson, L. D., Zavagno, A., Barlow, M. J., García-Lario, P., & Noriega-Crespo, A. 2012a, *A&A*, 537, A1
- Anderson, L. D., Zavagno, A., Deharveng, L., et al. 2012b, *A&A*, 542, A10
- Aniano, G., Draine, B. T., Gordon, K. D., & Sandstrom, K. 2011, *PASP*, 123, 1218
- Arthur, S. J., Kurtz, S. E., Franco, J., & Albarrán, M. Y. 2004, *ApJ*, 608, 282
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Becker, R. H., White, R. L., Helfand, D. J., & Zoonematkermani, S. 1994, *ApJS*, 91, 347
- Bendo, G. J., Draine, B. T., Engelbracht, C. W., et al. 2008, *MNRAS*, 389, 629
- Churchwell, E., Povich, M. S., Allen, D., et al. 2006, *ApJ*, 649, 759
- Churchwell, E., Watson, D. F., Povich, M. S., et al. 2007, *ApJ*, 670, 428
- Deharveng, L., Lefloch, B., Kurtz, S., et al. 2008, *A&A*, 482, 585
- Deharveng, L., & Zavagno, A. 2008, in *Astronomical Society of the Pacific Conference Series*, 387, *Massive Star Formation: Observations Confront Theory*, eds. H. Beuther, H. Linz, & T. Henning, 338
- Deharveng, L., Schuller, F., Anderson, L. D., et al. 2010, *A&A*, 523, A6
- Deharveng, L., Zavagno, A., Anderson, L. D., et al. 2012, *A&A*, 546, A74
- Deharveng, L., Zavagno, A., Samal, M. R., et al. 2015, *A&A*, 582, A1
- Draine, B. T. 2011a, *ApJ*, 732, 100
- Draine, B. T. 2011b, *Physics of the Interstellar and Intergalactic Medium* (Princeton: Princeton Univ. Press)
- Draine, B. T., & Li, A. 2007, *ApJ*, 657, 810
- Egan, M. P., Price, S. D., Kraemer, K. E., et al. 2003, *VizieR Online Data Catalog*, 5114
- Fazio, G. G., Hora, J. L., Allen, L. E., et al. 2004, *ApJS*, 154, 10
- Gail, H. P., & Sedlmayr, E. 1979, *A&A*, 77, 165
- Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, *A&A*, 518, L3
- Khrantsova, M. S., Wiebe, D. S., Boley, P. A., & Pavlyuchenkov, Y. N. 2013, *MNRAS*, 431, 2006
- Lebouteiller, V., Brandl, B., Bernard-Salas, J., Devost, D., & Houck, J. R. 2007, *ApJ*, 665, 390
- Madden, S. C., Galliano, F., Jones, A. P., & Sauvage, M. 2006, *A&A*, 446, 877
- Makai, Z., Anderson, L. D., Mascoop, J. L., & Johnstone, B. 2017, *ApJ*, 846, 64
- Martins, F., Pomarès, M., Deharveng, L., Zavagno, A., & Bouret, J. C. 2010, *A&A*, 510, A32
- O’Dell, C. R., & Yusef-Zadeh, F. 2000, *AJ*, 120, 382
- Paladini, R., Umana, G., Veneziani, M., et al. 2012, *ApJ*, 760, 149
- Pavlyuchenkov, Y. N., Kirsanova, M. S., & Wiebe, D. S. 2013, *Astronomy Reports*, 57, 573
- Poglitsch, A., Waelkens, C., Geis, N., et al. 2010, *A&A*, 518, L2
- Rieke, G. H., Young, E. T., Engelbracht, C. W., et al. 2004, *ApJS*, 154, 25
- Samus, N. N., & Li, Y., *RAA (Research in Astronomy and Astrophysics)*, 2018, 18, 88
- Simpson, R. J., Povich, M. S., Kendrew, S., et al. 2012, *MNRAS*, 424, 2442
- Topchieva, A. P., Wiebe, D. S., Kirsanova, M. S., & Krushinskii, V. V. 2017, *Astronomy Reports*, 61, 1015
- Urquhart, J. S., Hoare, M. G., Lumsden, S. L., et al. 2009, *A&A*, 507, 795
- van Buren, D., & McCray, R. 1988, *ApJ*, 329, L93
- Zinnecker, H., & Yorke, H. W. 2007, *ARA&A*, 45, 481